

Case Number:	CM15-0012379		
Date Assigned:	03/17/2015	Date of Injury:	07/09/2013
Decision Date:	04/16/2015	UR Denial Date:	01/07/2015
Priority:	Standard	Application Received:	01/21/2015

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:
 State(s) of Licensure: New York, Tennessee
 Certification(s)/Specialty: Emergency Medicine

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The injured worker is a 35 year old male, who sustained an industrial injury on 7/9/13. The injured worker has complaints of low back pain described as sharp, stabbing, burning and constant. The diagnoses have included low back pain; lumbar disc displacement and lumbar radiculopathy. The PR2 dated 1/8/15 noted that the injured worker has constant pain in the mid-low back radiating into the lower extremities. The pain is aggravated by bending, twisting, pushing, pulling, prolonged sitting, prolonged standing and walking multiple blocks.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

Lumbar Steroid Injection L4 (x1) with Epidurography: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Epidural Steroid Injections Page(s): 46.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Epidural Steroid Injections Page(s): 46.

Decision rationale: Epidural steroid injections are recommended as an option for treatment of radicular pain (defined as pain in dermatomal distribution with corroborative findings of radiculopathy). Radiculopathy must be documented by physical examination and corroborated by imaging studies and/or electrodiagnostic testing. Epidural steroid injection can offer short term pain relief and use should be in conjunction with other rehab efforts, including continuing a home exercise program. There is little information on improved function. The American Academy of Neurology recently concluded that epidural steroid injections may lead to an improvement in radicular lumbosacral pain between 2 and 6 weeks following the injection, but they do not affect impairment of function or the need for surgery and do not provide long-term pain relief beyond 3 months, and there is insufficient evidence to make any recommendation for the use of epidural steroid injections to treat radicular cervical pain. In this case there is no documentation of corroboration by imaging or electrodiagnostic studies. Criteria for epidural steroid injection have not been met. The request should not be authorized.

Monitored Anesthesia Care: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Epidural Steroid Injections Page(s): 46.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Epidural Steroid Injections Page(s): 46.

Decision rationale: The monitored anesthesia care is for administration of epidural steroid injection. Epidural steroid injections are recommended as an option for treatment of radicular pain (defined as pain in dermatomal distribution with corroborative findings of radiculopathy). Radiculopathy must be documented by physical examination and corroborated by imaging studies and/or electrodiagnostic testing. Epidural steroid injection can offer short term pain relief and use should be in conjunction with other rehab efforts, including continuing a home exercise program. There is little information on improved function. The American Academy of Neurology recently concluded that epidural steroid injections may lead to an improvement in radicular lumbosacral pain between 2 and 6 weeks following the injection, but they do not affect impairment of function or the need for surgery and do not provide long-term pain relief beyond 3 months, and there is insufficient evidence to make any recommendation for the use of epidural steroid injections to treat radicular cervical pain. In this case there is no documentation of corroboration by imaging or electrodiagnostic studies. Criteria for epidural steroid injection have not been met. Epidural steroid injection is not indicated. Monitored anesthesia care is not necessary. The request should not be authorized.