

Case Number:	CM15-0018519		
Date Assigned:	02/06/2015	Date of Injury:	01/31/2011
Decision Date:	04/07/2015	UR Denial Date:	01/03/2015
Priority:	Standard	Application Received:	01/31/2015

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:
 State(s) of Licensure: California, District of Columbia, Maryland
 Certification(s)/Specialty: Anesthesiology, Pain Management

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

A 34 year old female who sustained an industrial injury on 01/31/2011. Current diagnoses include left cervical strain-chronic, lumbosacral strain-chronic, carpal tunnel syndrome-chronic, right hip strain-chronic, chronic knee pain, and dyspepsia-mild. Previous treatments included medication management and TENS unit. Report dated 01/06/2015 noted that the injured worker presented with complaints that included neck, low back, right leg, and right upper extremity pain. Physical examination was positive for abnormal findings. Utilization review performed on 01/03/2015 non-certified a prescription for Protonix, based on clinical information submitted does not support medical necessity. The reviewer referenced the Official disability Guidelines in making this decision.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

Protonix 20mg tab #90: Upheld

Claims Administrator guideline: The Claims Administrator did not base their decision on the MTUS. Decision based on Non-MTUS Citation Official Disability Guidelines.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines NSAIDs, GI symptoms & cardiovascular risk Page(s): 68.

Decision rationale: In the treatment of dyspepsia secondary to NSAID therapy, the MTUS recommends stopping the NSAID, switching to a different NSAID, or considering the use of an H2-receptor antagonist or a PPI. The MTUS Chronic Pain Medical Treatment Guidelines recommend the use of proton pump inhibitors in conjunction with NSAIDs in situations in which the patient is at risk for gastrointestinal events including: (1) age > 65 years; (2) history of peptic ulcer, GI bleeding or perforation; (3) concurrent use of ASA, corticosteroids, and/or an anticoagulant; or (4) high dose/multiple NSAID (e.g., NSAID + low-dose ASA). CPMTG guidelines further specify: "Recommendations: Patients with no risk factor and no cardiovascular disease: Non-selective NSAIDs OK (e.g, ibuprofen, naproxen, etc.) Patients at intermediate risk for gastrointestinal events and no cardiovascular disease: (1) A non-selective NSAID with either a PPI (Proton Pump Inhibitor, for example, 20 mg omeprazole daily) or misoprostol (200g four times daily) or (2) a Cox-2 selective agent. Long-term PPI use (> 1 year) has been shown to increase the risk of hip fracture (adjusted odds ratio 1.44). Patients at high risk for gastrointestinal events with no cardiovascular disease: A Cox-2 selective agent plus a PPI if absolutely necessary. Patients at high risk of gastrointestinal events with cardiovascular disease: If GI risk is high the suggestion is for a low-dose Cox-2 plus low dose Aspirin (for cardioprotection) and a PPI. If cardiovascular risk is greater than GI risk the suggestion is naproxyn plus low-dose aspirin plus a PPI. (Laine, 2006) (Scholmerich, 2006) (Nielsen, 2006) (Chan, 2004) (Gold, 2007) (Laine, 2007)"As there is no documentation of peptic ulcer, GI bleeding or perforation, or cardiovascular disease in the records available for my review, the injured worker's risk for gastrointestinal events is low, as such, medical necessity cannot be affirmed. In 2014, the PTP documented that the IW had dyspepsia with NSAID use, and failed first-line omeprazole for this. I respectfully disagree with the UR physician's assertion that there is no documentation of GI risk.