

Case Number:	CM15-0130025		
Date Assigned:	07/16/2015	Date of Injury:	11/11/2014
Decision Date:	08/12/2015	UR Denial Date:	06/25/2015
Priority:	Standard	Application Received:	07/06/2015

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:
 State(s) of Licensure: New Jersey, Alabama, California
 Certification(s)/Specialty: Neurology, Neuromuscular Medicine

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

This 43 year old male sustained an industrial injury to the left hand on 11/11/14. X-ray of the left hand (11/13/14) revealed a fractured middle finger. Previous treatment included physical therapy, splinting and medications. The injured worker denied any improvement following physical therapy. The number of previous therapy sessions was unclear. In a Doctor's First Report of Occupational injury dated 6/1/15, the injured worker complained of pain, stiffness, swelling, numbness and tingling to the left middle, ring and small fingers. The injured worker rated his pain 5-9/10 on the visual analog scale. The injured worker reported having difficulty gripping and grasping. Left hand x-rays (6/1/15), showed a healed fracture of the volar base of the middle phalanx of the long finger with satisfactory alignment. Current diagnoses included left long finger middle phalanx fracture, left hand crush injury, severe contractures of the left hand long finger and ring finger. The physician recommended occupational therapy three times a week for six weeks to improve range of motion and function of the left hand.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

Occupational therapy 3 x 6, 3rd finger left: Upheld

Claims Administrator guideline: The Claims Administrator did not base their decision on the MTUS. Decision based on Non-MTUS Citation Official Disability Guidelines (ODG).

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Physical Medicine Page(s): 98.

Decision rationale: According to MTUS guidelines, Physical Medicine is "Recommended as indicated below. Passive therapy (those treatment modalities that do not require energy expenditure on the part of the patient) can provide short term relief during the early phases of pain treatment and are directed at controlling symptoms such as pain, inflammation and swelling and to improve the rate of healing soft tissue injuries. They can be used sparingly with active therapies to help control swelling, pain and inflammation during the rehabilitation process. Active therapy is based on the philosophy that therapeutic exercise and/or activity are beneficial for restoring flexibility, strength, endurance, function, range of motion, and can alleviate discomfort. Active therapy requires an internal effort by the individual to complete a specific exercise or task. This form of therapy may require supervision from a therapist or medical provider such as verbal, visual and/or tactile instruction(s). Patients are instructed and expected to continue active therapies at home as an extension of the treatment process in order to maintain improvement levels. Home exercise can include exercise with or without mechanical assistance or resistance and functional activities with assistive devices. (Colorado, 2002) (Airaksinen, 2006) Patient-specific hand therapy is very important in reducing swelling, decreasing pain, and improving range of motion in CRPS. (Li, 2005) The use of active treatment modalities (e.g., exercise, education, activity modification) instead of passive treatments is associated with substantially better clinical outcomes. In a large case series of patients with low back pain treated by physical therapists, those adhering to guidelines for active rather than passive treatments incurred fewer treatment visits, cost less, and had less pain and less disability. The overall success rates were 64.7% among those adhering to the active treatment recommendations versus 36.5% for passive treatment. (Fritz, 2007)." The patient underwent a certain number of physical therapy without clear evidence of efficacy. There is no documentation that the patient cannot perform home exercise. Therefore, the request for 18 occupational therapy sessions for the 3rd finger left is not medically necessary.