

<b>Case Number:</b>	CM15-0130699		
<b>Date Assigned:</b>	08/18/2015	<b>Date of Injury:</b>	05/27/2009
<b>Decision Date:</b>	09/14/2015	<b>UR Denial Date:</b>	06/17/2015
<b>Priority:</b>	Standard	<b>Application Received:</b>	07/07/2015

### HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:  
 State(s) of Licensure: North Carolina  
 Certification(s)/Specialty: Family Practice

### CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The injured worker is a 47 year old female, who sustained an industrial injury on May 27, 2009. The injured worker was diagnosed as having lumbosacral neuritis, chronic back pain, lumbar degenerative disc disease (DDD) and lumbar radiculitis. Treatment to date has included therapy, oral and topical medication and epidural steroid injection. A progress note dated June 5, 2015 provides the injured worker complains of low back pain radiating down leg to the foot with the right greater than the left. She reports prior epidural steroid injection provided relief to the point she did not need pain medication for 3 weeks. Physical exam notes lumbar tenderness to palpation with full range of motion (ROM). The plan includes oral and topical medication.

### IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

**Gabapentin 300mg, #60 with 1 refill:** Overturned

**Claims Administrator guideline:** Decision based on MTUS Chronic Pain Treatment Guidelines Antiepilepsy drugs (AEDs); Specific Anti-Epilepsy Drugs-Gabapentin (Neurontin, Gabarone, generic available).

**MAXIMUS guideline:** Decision based on MTUS Chronic Pain Treatment Guidelines gabapentin Page(s): 18.

**Decision rationale:** The California chronic pain medical treatment guidelines section on Neurontin states: Gabapentin (Neurontin, Gabarone, generic available) has been shown to be effective for treatment of diabetic painful neuropathy and post-herpetic neuralgia and has been considered as a first-line treatment for neuropathic pain. (Backonja, 2002) (ICSI, 2007) (Knotkova, 2007) (Eisenberg, 2007) (Attal, 2006) This RCT concluded that gabapentin monotherapy appears to be efficacious for the treatment of pain and sleep interference associated with diabetic peripheral neuropathy and exhibits positive effects on mood and quality of life. (Backonja, 1998) It has been given FDA approval for treatment of post-herpetic neuralgia. The number needed to treat (NNT) for overall neuropathic pain is 4. It has a more favorable side-effect profile than Carbamazepine, with a number needed to harm of 2.5. (Wiffen2-Cochrane, 2005) (Zaremba, 2006) Gabapentin in combination with morphine has been studied for treatment of diabetic neuropathy and post-herpetic neuralgia. When used in combination the maximum tolerated dosage of both drugs was lower than when each was used as a single agent and better analgesia occurred at lower doses of each. (Gilron-NEJM, 2005) Recommendations involving combination therapy require further study. The requested medication is a first line agent to treatment neuropathic pain. The patient does have a diagnosis of neuropathic pain in the form of lumbar radiculopathy. Therefore the request is medically indicated.

**Ibuprofen 600mg, #60 with 5 refills:** Overturned

**Claims Administrator guideline:** Decision based on MTUS Chronic Pain Treatment Guidelines NSAIDs (non-steroidal anti-inflammatory drugs) Back Pain-Chronic low back pain; NSAIDs, GI symptoms & cardiovascular risk.

**MAXIMUS guideline:** Decision based on MTUS Chronic Pain Treatment Guidelines NSAID Page(s): 68-72.

**Decision rationale:** The California chronic pain medical treatment guidelines section on NSAID therapy states: Recommended at the lowest dose for the shortest period in patients with moderate to severe pain. Acetaminophen may be considered for initial therapy for patients with mild to moderate pain, and in particular, for those with gastrointestinal, cardiovascular or renovascular risk factors. NSAIDs appear to be superior to acetaminophen, particularly for patients with moderate to severe pain. There is no evidence to recommend one drug in this class over another based on efficacy. In particular, there appears to be no difference between traditional NSAIDs and COX-2 NSAIDs in terms of pain relief. The main concern of selection is based on adverse effects. COX 2 NSAIDs have fewer GI side effects at the risk of increased cardiovascular side effects, although the FDA has concluded that long-term clinical trials are best interpreted to suggest that cardiovascular risk occurs with all NSAIDs and is a class effect (with naproxyn being the safest drug). There is no evidence of long-term effectiveness for pain or function. (Chen, 2008) (Laine, 2008) This medication is recommended for the shortest period of time and at the lowest dose possible. The shortest period of time is not defined in the California MTUS. The requested medication is within the maximum dosing guidelines per the California MTUS. Therefore the request is certified. Therefore, the requested treatment is not medically necessary.

**Lidocaine 5% topical ointment, 50gm with 2 refills:** Upheld

**Claims Administrator guideline:** Decision based on MTUS Chronic Pain Treatment Guidelines Topical Analgesics-Lidocaine Indication: Neuropathic pain.

**MAXIMUS guideline:** Decision based on MTUS Chronic Pain Treatment Guidelines topical lidocaine Page(s): 111-112.

**Decision rationale:** The California chronic pain medical treatment guidelines section on topical lidocaine states: Lidocaine Indication: Neuropathic pain Recommended for localized peripheral pain after there has been evidence of a trial of first-line therapy (tri-cyclic or SNRI anti-depressants or an AED such as gabapentin or Lyrica). Topical lidocaine, in the formulation of a dermal patch (Lidoderm) has been designated for orphan status by the FDA for neuropathic pain. Lidoderm is also used off-label for diabetic neuropathy. No other commercially approved topical formulations of lidocaine (whether creams, lotions or gels) are indicated for neuropathic pain. Non-dermal patch formulations are generally indicated as local anesthetics and anti-pruritics. Further research is needed to recommend this treatment for chronic neuropathic pain disorders other than post-herpetic neuralgia. Formulations that do not involve a dermal-patch system are generally indicated as local anesthetics and anti-pruritics. In February 2007 the FDA notified consumers and healthcare professionals of the potential hazards of the use of topical lidocaine. Those at particular risk were individuals that applied large amounts of this substance over large areas, left the products on for long periods of time, or used the agent with occlusive dressings. Systemic exposure was highly variable among patients. Only FDA-approved products are currently recommended. (Argoff, 2006) (Dworkin, 2007) (Khaliq-Cochrane, 2007) (Knotkova, 2007) (Lexi-Comp, 2008) Non-neuropathic pain: Not recommended. There is only one trial that tested 4% lidocaine for treatment of chronic muscle pain. The results showed there was no superiority over placebo. (Scudds, 1995) This medication is recommended for localized peripheral pain. There is no documentation of failure of first line neuropathic pain medications. Therefore criteria as set forth by the California MTUS as outlined above have not been met and the request is not certified. Therefore, the requested treatment is not medically necessary.