

Case Number:	CM15-0131892		
Date Assigned:	07/20/2015	Date of Injury:	01/08/2015
Decision Date:	08/13/2015	UR Denial Date:	06/30/2015
Priority:	Standard	Application Received:	07/08/2015

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:

State(s) of Licensure: Iowa, Illinois, Hawaii

Certification(s)/Specialty: Preventive Medicine, Occupational Medicine, Public Health & General Preventive Medicine

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The injured worker (IW) is a 30 year old male who sustained an industrial injury on 01/08/2015. He reported pain in his left wrist and shoulder after unloading an item weighing about 55-65 lbs. The injured worker was diagnosed as having left lateral epicondylitis, left hand weakness with intermittent paresthesias, left shoulder weakness and impingement. Treatment to date has included medications for pain. Currently, the injured worker complains of numbness and tingling in the left shoulder, left arm, and all five digits of the left hand. Pain in the left shoulder blade extends to the 1 left arm and all five fingers of the left hand. He has constant weakness and fatigue in the left arm and swelling of the left hand. The symptoms are increased with activity and reach an 8 on a scale of 1-10. On examination, the worker exhibits 4/5 strength throughout the left arm an electromyogram shows bilateral mild carpal tunnel syndrome, and no evidence of cervical radiculopathy, brachial plexopathy, or other peripheral nerve entrapment (06/30/2015). He receives Tramadol for pain. A request for authorization was made for a left elbow MRI.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

Left elbow MRI: Upheld

Claims Administrator guideline: The Claims Administrator did not cite any medical evidence for its decision.

MAXIMUS guideline: Decision based on MTUS ACOEM Chapter 10 Elbow Disorders (Revised 2007) Page(s): 33-34. Decision based on Non-MTUS Citation Official Disability Guidelines (ODG) Elbow (Acute & Chronic), MRIs.

Decision rationale: ACOEM states, Criteria for ordering imaging studies are: The imaging study results will substantially change the treatment plan. Emergence of a red flag. Failure to progress in a rehabilitation program, evidence of significant tissue insult or neurological dysfunction that has been shown to be correctible by invasive treatment, and agreement by the patient to undergo invasive treatment if the presence of the correctible lesion is confirmed. For most patients presenting with elbow problems, special studies are not needed unless a period of at least 4 weeks of conservative care and observation fails to improve their symptoms. Most patients improve quickly, provided red flag conditions are ruled out. There are a few exceptions to the rule to avoid special studies absent red flags in the first month. These exceptions include: Plain-film radiography to rule out osteomyelitis or joint effusion in cases of significant septic olecranon bursitis. Electromyography (EMG) study if cervical radiculopathy is suspected as a cause of lateral arm pain, and that condition has been present for at least 6 weeks. Nerve conduction study and possibly EMG if severe nerve entrapment is suspected on the basis of physical examination, denervation atrophy is likely, and there is a failure to respond to conservative treatment. For patients with limitations of activity after 4 weeks and unexplained physical findings such as effusion or localized pain (especially following exercise), imaging may be indicated to clarify the diagnosis and revise the treatment strategy if appropriate. Imaging findings should be correlated with physical findings. In general, an imaging study may be an appropriate consideration for a patient whose limitations due to consistent symptoms have persisted for 1 month or more, as in the following cases: When surgery is being considered for a specific anatomic defect. To further evaluate potentially serious pathology, such as a possible tumor, when the clinical examination suggests the diagnosis. ACOEM further recommends MRI for suspected ulnar collateral ligament tears and recommends against MRI for suspected epicondylgia. ODG writes regarding elbow MRI, "Recommended as indicated below. Magnetic resonance imaging may provide important diagnostic information for evaluating the adult elbow in many different conditions, including: collateral ligament injury, epicondylitis, injury to the biceps and triceps tendons, abnormality of the ulnar, radial, or median nerve, and for masses about the elbow joint. There is a lack of studies showing the sensitivity and specificity of MR in many of these entities; most of the studies demonstrate MR findings in patients either known or highly likely to have a specific condition. Epicondylitis (lateral - "tennis elbow" or medial - in pitchers, golfers, and tennis players) is a common clinical diagnosis, and MRI is usually not necessary. Magnetic resonance may be useful for confirmation of the diagnosis in refractory cases and to exclude associated tendon and ligament tear. Indications for imaging -- Magnetic resonance imaging (MRI): Chronic elbow pain, suspect intra-articular osteocartilaginous body; plain films nondiagnostic, Chronic elbow pain, suspect occult injury; e.g., osteochondral injury; plain films--nondiagnostic, Chronic elbow pain, suspect unstable osteochondral injury; plain films nondiagnostic, Chronic elbow pain, suspect nerve entrapment or mass; plain films nondiagnostic, Chronic elbow pain, suspect chronic epicondylitis; plain films nondiagnostic, Chronic elbow pain, suspect collateral ligament tear; plain films nondiagnostic, Chronic elbow

pain, suspect biceps tendon tear and/or bursitis; plain films nondiagnostic. Repeat MRI is not routinely recommended, and should be reserved for a significant change in symptoms and/or findings suggestive of significant pathology. The medical notes provided did not document (physical exam, objective testing, or subjective complaints) any red flags, significant worsening in symptoms or other findings suggestive of the pathologies outlined in the above guidelines. The treatment notes do not indicate other extenuating circumstances to warrant deviation from the guidelines. As such, the request for Left elbow MRI is not medically necessary.