MAXIMUS Federal Services

Case Number:	CM15-0136307		
Date Assigned:	07/24/2015	Date of Injury:	02/16/2012
Decision Date:	08/27/2015	UR Denial Date:	07/06/2015
Priority:	Standard	Application Received:	07/14/2015

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:
State(s) of Licensure: California, District of Columbia, Maryland
Certification(s)/Specialty: Anesthesiology, Pain Management

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The injured worker is a 66 year old male, who sustained an industrial injury on 2/16/2012. Diagnoses include rotator cuff syndrome shoulder and neck sprain/strain. Treatment to date has included conservative measures including medication management. Current medications include Tramadol and LidoPro. Per the Primary Treating Physician's Progress Report dated 6/05/2015, the injured worker reported worsening pain in the left side of his neck and pain from his left shoulder radiating into his cervical spine and upper arm. Physical examination revealed significant tenderness in the left splenius capitis and cervical paravertebral region. The plan of care included physical therapy and oral and topical medications and authorization was requested for LidoPro topical ointment.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

Compound medication: LidoPro topical ointment #2: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Topical Analgesics Page(s): 60, 111-113.

Decision rationale: LidoPro contains capsaicin, lidocaine, menthol, methyl salicylate. Per MTUS p 112 with regard to capsaicin, "Indications: There are positive randomized studies with capsaicin cream in patients with osteoarthritis, fibromyalgia, and chronic non-specific back pain, but it should be considered experimental in very high doses. Although topical capsaicin has moderate to poor efficacy, it may be particularly useful (alone or in conjunction with other modalities) in patients whose pain has not been controlled successfully with conventional therapy." Methyl salicylate may have an indication for chronic pain in this context. Per MTUS p105, "Recommended. Topical salicylate (e.g., Ben-Gay, methyl salicylate) is significantly better than placebo in chronic pain. (Mason-BMJ, 2004)." However, the other ingredients in LidoPro are not indicated. The preponderance of evidence indicates that overall this medication is not medically necessary. Regarding topical lidocaine, MTUS states (p112) "Neuropathic pain Recommended for localized peripheral pain after there has been evidence of a trial of first-line therapy (tri-cyclic or SNRI anti-depressants or an AED such as gabapentin or Lyrica). Non-neuropathic pain: Not recommended. There is only one trial that tested 4% lidocaine for treatment of chronic muscle pain. The results showed there was no superiority over placebo. (Scudds, 1995) “The documentation submitted for review does not contain evidence of trial of first-line therapy to support the use of topical lidocaine. LidoPro topical lotion contains menthol. The CA MTUS, ODG, National Guidelines Clearinghouse, and ACOEM provide no evidence-based recommendations regarding the topical application of menthol. It is the opinion of this IMR reviewer that a lack of endorsement, a lack of mention, inherently implies a lack of recommendation, or a status equivalent to "not recommended." Since menthol is not medically indicated, then the overall product is not indicated per MTUS as outlined below. Note the statement on page 111: Any compounded product that contains at least one drug (or drug class) that is not recommended is not recommended. Regarding the use of multiple medications, MTUS p60 states "only one medication should be given at a time, and interventions that are active and passive should remain unchanged at the time of the medication change. A trial should be given for each individual medication. Analgesic medications should show effects within 1 to 3 days, and the analgesic effect of antidepressants should occur within 1 week. A record of pain and function with the medication should be recorded. (Mens, 2005) The recent AHRQ review of comparative effectiveness and safety of analgesics for osteoarthritis concluded that each of the analgesics was associated with a unique set of benefits and risks, and no currently available analgesic was identified as offering a clear overall advantage compared with the others." Therefore, it would be optimal to trial each medication individually.