

Case Number:	CM15-0136375		
Date Assigned:	07/24/2015	Date of Injury:	08/20/2013
Decision Date:	08/21/2015	UR Denial Date:	06/23/2015
Priority:	Standard	Application Received:	07/14/2015

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:
 State(s) of Licensure: North Carolina
 Certification(s)/Specialty: Family Practice

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The injured worker is a (n) 45-year-old male, who sustained an industrial injury on 8/20/13. He reported injury to his left knee related to a fall. The injured worker was diagnosed as having derangement meniscus, muscle atrophy of the left quad and chronic pain syndrome. Treatment to date has included physical therapy, a left knee MRI showing a meniscus tear in 2013, psychiatric treatments and Norco. As of the PR2 dated 3/23/15, the injured worker reports left knee pain, gait instability, depression and increased pain with walking. Objective findings include moderate left knee swelling and painful flexion. The treating physician requested 8 sessions of transitional step down program 5 hours per session 1 per week for 8 weeks and physical therapy 2 x weekly for 8 weeks.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

8 sessions of transitional step down program 5 hours per session 1 per week for 8 weeks:
 Upheld

Claims Administrator guideline: The Claims Administrator did not cite any medical evidence for its decision.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines functional restoration program Page(s): 49.

Decision rationale: The California chronic pain medical treatment guidelines section on functional restoration programs states: Recommended, although research is still ongoing as to how to most appropriately screen for inclusion in these programs. Functional restoration programs (FRPs), a type of treatment included in the category of interdisciplinary pain programs (see chronic pain programs), were originally developed by Mayer and Gatchel. FRPs were designed to use a medically directed, interdisciplinary pain management approach geared specifically to patients with chronic disabling occupational musculoskeletal disorders. These programs emphasize the importance of function over the elimination of pain. FRPs incorporate components of exercise progression with disability management and psychosocial intervention. Long-term evidence suggests that the benefit of these programs diminishes over time, but remains positive when compared to cohorts that did not receive an intensive program. (Bendix, 1998) A Cochrane review suggests that there is strong evidence that intensive multidisciplinary rehabilitation with functional restoration reduces pain and improves function of patients with low back pain. The evidence is contradictory when evaluating the programs in terms of vocational outcomes. (Guzman 2001) It must be noted that all studies used for the Cochrane review excluded individuals with extensive radiculopathy, and several of the studies excluded patients who were receiving a pension, limiting the generalizability of the above results. Studies published after the Cochrane review also indicate that intensive programs show greater effectiveness, in particular in terms of return to work, than less intensive treatment. (Airaksinen, 2006) There appears to be little scientific evidence for the effectiveness of multidisciplinary biopsychosocial rehabilitation compared with other rehabilitation facilities for neck, shoulder pain, as opposed to low back pain, and generalized pain syndromes. (Karjalainen, 2003) Treatment is not suggested for longer than 2 weeks without evidence of demonstrated efficacy as documented by subjective and objective gains. For general information, see Chronic pain programs. While functional restoration programs are recommended per the California MTUS, the length of time is for 2 weeks unless there is documentation of demonstrated efficacy by subjective and objective gains. The patient has completed the recommended amount of time for this program. There is no objective documented need for a step down program and the request is not medically necessary.

Physical therapy 2 times a week for 8 weeks: Upheld

Claims Administrator guideline: The Claims Administrator did not cite any medical evidence for its decision.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines physical medicine Page(s): 98-99.

Decision rationale: The California chronic pain medical treatment guidelines section on physical medicine states: Recommended as indicated below. Passive therapy (those treatment modalities that do not require energy expenditure on the part of the patient) can provide short term relief during the early phases of pain treatment and are directed at controlling symptoms such as pain, inflammation and swelling and to improve the rate of healing soft tissue injuries.

They can be used sparingly with active therapies to help control swelling, pain and inflammation during the rehabilitation process. Active therapy is based on the philosophy that therapeutic exercise and/or activity are beneficial for restoring flexibility, strength, endurance, function, range of motion, and can alleviate discomfort. Active therapy requires an internal effort by the individual to complete a specific exercise or task. This form of therapy may require supervision from a therapist or medical provider such as verbal, visual and/or tactile instruction(s). Patients are instructed and expected to continue active therapies at home as an extension of the treatment process in order to maintain improvement levels. Home exercise can include exercise with or without mechanical assistance or resistance and functional activities with assistive devices. (Colorado, 2002) (Airaksinen, 2006) Patient-specific hand therapy is very important in reducing swelling, decreasing pain, and improving range of motion in CRPS. (Li, 2005) The use of active treatment modalities (e.g., exercise, education, activity modification) instead of passive treatments is associated with substantially better clinical outcomes. In a large case series of patients with low back pain treated by physical therapists, those adhering to guidelines for active rather than passive treatments incurred fewer treatment visits, cost less, and had less pain and less disability. The overall success rates were 64.7% among those adhering to the active treatment recommendations versus 36.5% for passive treatment. (Fritz, 2007) Physical Medicine Guidelines; Allow for fading of treatment frequency (from up to 3 visits per week to 1 or less), plus active self-directed home Physical Medicine. Myalgia and myositis, unspecified (ICD9 729.1): 9-10 visits over 8 weeks. Neuralgia, neuritis, and radiculitis, unspecified (ICD9 729.2) 8-10 visits over 4 weeks. Reflex sympathetic dystrophy (CRPS) (ICD9 337.2): 24 visits over 16 weeks. The goal of physical therapy is graduation to home therapy after a certain amount of recommended sessions. The patient has already completed physical therapy. The request is in excess of these recommendations per the California MTUS. There is no objective reason why the patient would not be moved to home therapy after completing the recommended amount of supervised sessions in the provided clinical documentation. Therefore, the request is not medically necessary.