

Case Number:	CM15-0137557		
Date Assigned:	07/27/2015	Date of Injury:	07/25/2012
Decision Date:	08/24/2015	UR Denial Date:	07/06/2015
Priority:	Standard	Application Received:	07/15/2015

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:
 State(s) of Licensure: North Carolina
 Certification(s)/Specialty: Family Practice

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The injured worker is a 22 year old female who sustained an industrial/work injury on 7/25/12. She reported an initial complaint of lower back and leg pain. The injured worker was diagnosed as having lumbar radiculopathy, lumbosacral arthrosis, herniated disc of lumbar spine, and chronic pain syndrome. Treatment to date includes medication, VP shunt placement with long term chronic migraines, chiropractic treatments, and acupuncture. MRI results of the lumbar spine impression noted 2-3 mm posterior disc protrusion at L5-S1 which abuts the traversing left S1 nerve root with no stenosis found. Currently, the injured worker complained of low back, posterior thighs, and calves. Pain is described as sharp and throbbing and rated 7/10. Per the primary physician's report (PR-2) on 6/26/15, exam noted tenderness on palpation of bilateral lower paraspinals at L5-S1 levels, lumbospinal motion was severely limited with range of motion, and normal reflexes. The requested treatments include bilateral L5-S1 transforaminal epidural steroid.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

Bilateral L5-S1 transforaminal epidural steroid injection: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Epidural steroid injections Page(s): 46.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines epidural steroid injections Page(s): 46.

Decision rationale: The California chronic pain medical treatment guidelines section on epidural steroid injections (ESI) states: Criteria for the use of Epidural steroid injections: Note: The purpose of ESI is to reduce pain and inflammation, restoring range of motion and thereby facilitating progress in more active treatment programs, and avoiding surgery, but this treatment alone offers no significant long-term functional benefit. 1) Radiculopathy must be documented by physical examination and corroborated by imaging studies and/or electrodiagnostic testing. 2) Initially unresponsive to conservative treatment (exercises, physical methods, NSAIDs and muscle relaxants). 3) Injections should be performed using fluoroscopy (live x-ray) for guidance. 4) If used for diagnostic purposes, a maximum of two injections should be performed. A second block is not recommended if there is inadequate response to the first block. Diagnostic blocks should be at an interval of at least one to two weeks between injections. 5) No more than two nerve root levels should be injected using transforaminal blocks. 6) No more than one interlaminar level should be injected at one session. 7) In the therapeutic phase, repeat blocks should be based on continued objective documented pain and functional improvement, including at least 50% pain relief with associated reduction of medication use for six to eight weeks, with a general recommendation of no more than 4 blocks per region per year. (Manchikanti, 2003) (CMS, 2004) (Boswell, 2007) 8) Current research does not support a series-of-three injections in either the diagnostic or therapeutic phase. We recommend no more than 2 ESI injections. The patient has the documentation of low back pain however there is no included imaging or nerve conduction studies in the clinical documentation provided for review that collaborates dermatomal radiculopathy on exam for the requested level of ESI. Therefore, criteria have not been met and the request is not medically necessary.

Physical therapy 3 x 4 weeks: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Physical medicine Page(s): 98-99.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines physical medicine Page(s): 98-99.

Decision rationale: The California chronic pain medical treatment guidelines section on physical medicine states: Recommended as indicated below. Passive therapy (those treatment modalities that do not require energy expenditure on the part of the patient) can provide short term relief during the early phases of pain treatment and are directed at controlling symptoms such as pain, inflammation and swelling and to improve the rate of healing soft tissue injuries. They can be used sparingly with active therapies to help control swelling, pain and inflammation during the rehabilitation process. Active therapy is based on the philosophy that therapeutic exercise and/or activity are beneficial for restoring flexibility, strength, endurance, function, range of motion, and can alleviate discomfort. Active therapy requires an internal

effort by the individual to complete a specific exercise or task. This form of therapy may require supervision from a therapist or medical provider such as verbal, visual and/or tactile instruction(s). Patients are instructed and expected to continue active therapies at home as an extension of the treatment process in order to maintain improvement levels. Home exercise can include exercise with or without mechanical assistance or resistance and functional activities with assistive devices. (Colorado, 2002) (Airaksinen, 2006) Patient-specific hand therapy is very important in reducing swelling, decreasing pain, and improving range of motion in CRPS. (Li, 2005) The use of active treatment modalities (e.g., exercise, education, activity modification) instead of passive treatments is associated with substantially better clinical outcomes. In a large case series of patients with low back pain treated by physical therapists, those adhering to guidelines for active rather than passive treatments incurred fewer treatment visits, cost less, and had less pain and less disability. The overall success rates were 64.7% among those adhering to the active treatment recommendations versus 36.5% for passive treatment. (Fritz, 2007) Physical Medicine Guidelines: Allow for fading of treatment frequency (from up to 3 visits per week to 1 or less), plus active self-directed home Physical Medicine. Myalgia and myositis, unspecified (ICD9 729.1): 9-10 visits over 8 weeks. Neuralgia, neuritis, and radiculitis, unspecified (ICD9 729.2): 8-10 visits over 4 weeks. Reflex sympathetic dystrophy (CRPS) (ICD9 337.2): 24 visits over 16 weeks. The requested amount of physical therapy is in excess of California chronic pain medical treatment guidelines. The patient has already completed a course of physical therapy. There is no objective explanation why the patient would need excess physical therapy and not be transitioned to active self-directed physical medicine. The request is not medically necessary.