

Case Number:	CM15-0139564		
Date Assigned:	07/29/2015	Date of Injury:	07/26/2002
Decision Date:	09/01/2015	UR Denial Date:	06/29/2015
Priority:	Standard	Application Received:	07/20/2015

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:

State(s) of Licensure: Maryland, Texas, Virginia

Certification(s)/Specialty: Internal Medicine, Allergy and Immunology, Rheumatology

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The injured worker is a 55 year old male who sustained an industrial injury on 7/26/02. He had complaints of low back pain with radiation into the right lower extremity. Treatment includes: medication, physical therapy and injections. Office visit report dated 5/28/15 reports follow up for refill of current medications. He has been compliant for use of medications. Prescriptions: pantoprazole 20 mg 1 twice per day, #60 and Lunesta 3 mg 1 at night, #30. Diagnoses include: degeneration lumbar, disorders of sacrum, sprain/strain lumbar region and major depressive disorder. Work status: permanent and stationary. Office visit report dated 6/9/15 reports follow up post lumbar epidural injection. He has a reduction of his pain, especially with the pain that radiates down the right lower extremity. Prolonged sitting and standing still aggravates the pain. Plan of care: continue to monitor improvement with the epidural injection and medications refilled. Follow up in 4 weeks.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

Pantoprazole 20mg quantity: 60 with 3 refills: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines NSAIDs, GI symptoms & cardiovascular risk Page(s): 68-69. Decision based on Non-MTUS Citation Official Disability Guidelines (ODG) Pain (Chronic), NSAIDs, GI symptoms & cardiovascular risk.

Decision rationale: Protonix is the brand name version of Pantoprazole, which is a proton pump inhibitor. MTUS states, "Determine if the patient is at risk for gastrointestinal events: (1) age > 65 years; (2) history of peptic ulcer, GI bleeding or perforation; (3) concurrent use of ASA, corticosteroids, and/or an anticoagulant; or (4) high dose/multiple NSAID (e.g., NSAID + low-dose ASA)." And "Patients at intermediate risk for gastrointestinal events and no cardiovascular disease: (1) A non-selective NSAID with either a PPI (Proton Pump Inhibitor, for example, 20 mg omeprazole daily) or misoprostol (200g four times daily) or (2) a Cox-2 selective agent. Long-term PPI use (> 1 year) has been shown to increase the risk of hip fracture (adjusted odds ratio 1.44)." ODG states, "If a PPI is used, omeprazole OTC tablets or lansoprazole 24HR OTC are recommended for an equivalent clinical efficacy and significant cost savings. Products in this drug class have demonstrated equivalent clinical efficacy and safety at comparable doses, including esomeprazole (Nexium), lansoprazole (Prevacid), omeprazole (Prilosec), pantoprazole (Protonix), dexlansoprazole (Dexilant), and rabeprazole (Aciphex). (Shi, 2008) A trial of omeprazole or lansoprazole is recommended before Nexium therapy. The other PPIs, Protonix, Dexilant, and Aciphex, should also be second-line. According to the latest AHRQ Comparative Effectiveness Research, all of the commercially available PPIs appeared to be similarly effective. (AHRQ, 2011)". The patient does not meet the age recommendations for increased GI risk. The medical documents provided establish the patient had an Upper endoscopy but fail to demonstrate the results, with possible GERD allergies or esophagitis but does not indicate history of peptic ulcer, GI bleeding or perforation. Medical records do not indicate that the patient is on ASA, corticosteroids, and/or an anticoagulant; or high dose/multiple NSAID. Additionally per guidelines, Pantoprazole is considered second line therapy and the treating physician has not provided detailed documentation of a failed trial of omeprazole and/or lansoprazole. As such, the request for Pantoprazole 20mg quantity; 60 with 3 refills is not medically necessary.

Lunesta 3mg quantity: 30: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines.

MAXIMUS guideline: The Expert Reviewer did not base their decision on the MTUS. Decision based on Non-MTUS Citation Official Disability Guidelines (ODG) Chronic Pain, insomnia, Mental Illness, Eszopicolone (Lunesta).

Decision rationale: MTUS is silent specifically regarding eszopicolone (Lunesta), therefore other guidelines were utilized. ODG states regarding Eszopicolone, "Not recommended for long-term use, but recommended for short-term use. See Insomnia treatment. See also the Pain Chapter. Recommend limiting use of hypnotics to three weeks maximum in the first two months of injury only, and discourage use in the chronic phase". For insomnia ODG recommends that

"Pharmacological agents should only be used after careful evaluation of potential causes of sleep disturbance. Failure of sleep disturbance to resolve in a 7 to 10 day period may indicate a psychiatric and/or medical illness. (Lexi-Comp, 2008) Primary insomnia is generally addressed pharmacologically. Secondary insomnia may be treated with pharmacological and/or psychological measures. The specific component of insomnia should be addressed: (a) Sleep onset; (b) Sleep maintenance; (c) Sleep quality; & (d) Next-day functioning". Medical records do not indicate patient's sleep hygiene or the need for variance from the guidelines, such as "a) Wake at the same time everyday; (b) Maintain a consistent bedtime; (c) Exercise regularly (not within 2 to 4 hours of bedtime); (d) Perform relaxing activities before bedtime; (e) Keep your bedroom quiet and cool; (f) Do not watch the clock; (g) Avoid caffeine and nicotine for at least six hours before bed; (h) Only drink in moderation; & (i) Avoid napping". Medical documents indicate that the patient has been on Eszopicolone in excess of the guidelines. Additionally, medical records do not indicate what components of insomnia have been addressed, treated with conservative measures, and the results of those conservative treatments. As such, the request for 1 Lunesta 3mg quantity: 30 is not medically necessary.