

Case Number:	CM15-0139618		
Date Assigned:	07/29/2015	Date of Injury:	06/01/2005
Decision Date:	09/23/2015	UR Denial Date:	07/10/2015
Priority:	Standard	Application Received:	07/20/2015

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:

State(s) of Licensure: Oregon

Certification(s)/Specialty: Plastic Surgery, Hand Surgery

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

This 57 year old female sustained an industrial injury on 6/01/05. She subsequently reported wrist and hand pain. Diagnoses include left wrist radiocarpal arthritis, history of left carpal tunnel release and right thumb CMC joint arthritis. Treatments to date include x-ray testing, injections, and carpal tunnel surgery and prescription pain medications. The injured worker continues to experience left wrist pain. Upon examination, there was tenderness to touch over the radiocarpal joint dorsally. There was swelling noted. A request for Right thumb basal joint reconstruction and MCP volar capsulodesis, Pre-operative medical clearance: Electrocardiogram (EKG), Pre-operative medical clearance: Complete blood count (CBC) with auto diff wbc and Pre-operative medical clearance: Basic metabolic panel was made by the treating physician.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

Right thumb basal joint reconstruction and MCP volar capsulodesis: Overturned

Claims Administrator guideline: Decision based on MTUS ACOEM Chapter 11 Forearm, Wrist, and Hand Complaints Page(s): 270.

MAXIMUS guideline: The Expert Reviewer did not base their decision on the MTUS. Decision based on Non-MTUS Citation Official Disability Guidelines (ODG) Forearm, Wrist and Hand.

Decision rationale: According to the ODG guidelines with reference to CMC joint thumb arthroplasty, "In our series, total joint arthroplasty of the thumb CMC joint has proven to be efficacious with improved motion, strength, and pain relief for the treatment of stage III and early stage IV osteoarthritis of the CMC joint in older patients with low activity demands." The ODG guidelines support medical necessity for the proposed CMC joint replacement. This patient has advanced degenerative changes on X-ray and has persistent symptoms despite splinting and several steroid injections.

Pre-operative medical clearance: Electrocardiogram (EKG): Overturned

Claims Administrator guideline: The Claims Administrator did not cite any medical evidence for its decision.

MAXIMUS guideline: The Expert Reviewer did not base their decision on the MTUS. Decision based on Non-MTUS Citation ODG-TWC, Low Back updated 5/15/15.

Decision rationale: ODG-TWC, Low Back updated 5/15/15 states: "Preoperative testing (e.g., chest radiography, electrocardiography, laboratory testing, urinalysis) is often performed before surgical procedures. These investigations can be helpful to stratify risk, direct anesthetic choices, and guide postoperative management, but often are obtained because of protocol rather than medical necessity. The decision to order preoperative tests should be guided by the patient's clinical history, comorbidities, and physical examination findings. Patients with signs or symptoms of active cardiovascular disease should be evaluated with appropriate testing, regardless of their preoperative status. Electrocardiography is recommended for patients undergoing high-risk surgery and those undergoing intermediate-risk surgery who have additional risk factors. Patients undergoing low-risk surgery do not require electrocardiography. Chest radiography is reasonable for patients at risk of postoperative pulmonary complications if the results would change perioperative management. Patients in their usual state of health who are undergoing cataract surgery do not require preoperative testing. (Feely, 2013) Routine preoperative tests are defined as those done in the absence of any specific clinical indication or purpose and typically include a panel of blood tests, urine tests, chest radiography, and an electrocardiogram (ECG). These tests are performed to find latent abnormalities, such as anemia or silent heart disease, that could impact how, when, or whether the planned surgical procedure and concomitant anesthesia are performed. It is unclear whether the benefits accrued from responses to true-positive tests outweigh the harms of false-positive preoperative tests and, if there is a net benefit, how this benefit compares to the resource utilization required for testing. An alternative to routine preoperative testing for the purpose of determining fitness for anesthesia and identifying patients at high risk of postoperative complications may be to conduct a history and physical examination, with selective testing based on the clinician's findings. However, the relative effect on patient and surgical outcomes, as well as resource utilization, of these two approaches is unknown. (AHRQ, 2013) The latest AHRQ comparative effectiveness research on the benefits and harms of routine preoperative testing concludes that, except for

cataract surgery, there is insufficient evidence comparing routine and per-protocol testing." This patient has a history of obesity, COPD and acid reflux. She is over age 50. She will need general anesthesia. EKG is indicated.

Pre-operative medical clearance: Complete blood count (CBC) with auto diff wbc: Overturned

Claims Administrator guideline: The Claims Administrator did not cite any medical evidence for its decision.

MAXIMUS guideline: The Expert Reviewer did not base their decision on the MTUS. Decision based on Non-MTUS Citation ODG-TWC, Low Back updated 5/15/15.

Decision rationale: ODG-TWC, Low Back updated 5/15/15 states: "Preoperative testing (e.g., chest radiography, electrocardiography, laboratory testing, urinalysis) is often performed before surgical procedures. These investigations can be helpful to stratify risk, direct anesthetic choices, and guide postoperative management, but often are obtained because of protocol rather than medical necessity. The decision to order preoperative tests should be guided by the patient's clinical history, comorbidities, and physical examination findings. Patients with signs or symptoms of active cardiovascular disease should be evaluated with appropriate testing, regardless of their preoperative status. Electrocardiography is recommended for patients undergoing high-risk surgery and that undergoing intermediate-risk surgery who have additional risk factors. Patients undergoing low-risk surgery do not require electrocardiography. Chest radiography is reasonable for patients at risk of postoperative pulmonary complications if the results would change perioperative management. Patients in their usual state of health who are undergoing cataract surgery do not require preoperative testing. (Feely, 2013) Routine preoperative tests are defined as those done in the absence of any specific clinical indication or purpose and typically include a panel of blood tests, urine tests, chest radiography, and an electrocardiogram (ECG). These tests are performed to find latent abnormalities, such as anemia or silent heart disease, that could impact how, when, or whether the planned surgical procedure and concomitant anesthesia are performed. It is unclear whether the benefits accrued from responses to true-positive tests outweigh the harms of false-positive preoperative tests and, if there is a net benefit, how this benefit compares to the resource utilization required for testing. An alternative to routine preoperative testing for the purpose of determining fitness for anesthesia and identifying patients at high risk of postoperative complications may be to conduct a history and physical examination, with selective testing based on the clinician's findings. However, the relative effect on patient and surgical outcomes, as well as resource utilization, of these two approaches is unknown. (AHRQ, 2013) The latest AHRQ comparative effectiveness research on the benefits and harms of routine preoperative testing, concludes that, except for cataract surgery, there is insufficient evidence comparing routine and per-protocol testing." This patient has a history of obesity, COPD and acid reflux. She is over age 50. She will need general anesthesia. A preoperative with CBC and basic metabolic panel is indicated because of the increased likelihood of abnormalities due to her co-morbid conditions and medications.

Pre-operative medical clearance: Basic metabolic panel: Overturned

Claims Administrator guideline: The Claims Administrator did not cite any medical evidence for its decision.

MAXIMUS guideline: The Expert Reviewer did not base their decision on the MTUS. Decision based on Non-MTUS Citation ODG-TWC, Low Back updated 5/15/15.

Decision rationale: ODG-TWC, Low Back updated 5/15/15 states: "Preoperative testing (e.g., chest radiography, electrocardiography, laboratory testing, urinalysis) is often performed before surgical procedures. These investigations can be helpful to stratify risk, direct anesthetic choices, and guide postoperative management, but often are obtained because of protocol rather than medical necessity. The decision to order preoperative tests should be guided by the patient's clinical history, comorbidities, and physical examination findings. Patients with signs or symptoms of active cardiovascular disease should be evaluated with appropriate testing, regardless of their preoperative status. Electrocardiography is recommended for patients undergoing high-risk surgery and those undergoing intermediate-risk surgery who have additional risk factors. Patients undergoing low-risk surgery do not require electrocardiography. Chest radiography is reasonable for patients at risk of postoperative pulmonary complications if the results would change perioperative management. Patients in their usual state of health who are undergoing cataract surgery do not require preoperative testing. (Feely, 2013) Routine preoperative tests are defined as those done in the absence of any specific clinical indication or purpose and typically include a panel of blood tests, urine tests, chest radiography, and an electrocardiogram (ECG). These tests are performed to find latent abnormalities, such as anemia or silent heart disease, that could impact how, when, or whether the planned surgical procedure and concomitant anesthesia are performed. It is unclear whether the benefits accrued from responses to true-positive tests outweigh the harms of false-positive preoperative tests and, if there is a net benefit, how this benefit compares to the resource utilization required for testing. An alternative to routine preoperative testing for the purpose of determining fitness for anesthesia and identifying patients at high risk of postoperative complications may be to conduct a history and physical examination, with selective testing based on the clinician's findings. However, the relative effect on patient and surgical outcomes, as well as resource utilization, of these two approaches is unknown. (AHRQ, 2013) The latest AHRQ comparative effectiveness research on the benefits and harms of routine preoperative testing, concludes that, except for cataract surgery, there is insufficient evidence comparing routine and per-protocol testing." This patient has a history of obesity, COPD and acid reflux. She is over age 50. She will need general anesthesia. A preoperative with CBC and basic metabolic panel is indicated because of the increased likelihood of abnormalities due to her co-morbid conditions and medications.