

Case Number:	CM15-0140882		
Date Assigned:	07/30/2015	Date of Injury:	08/22/2011
Decision Date:	08/28/2015	UR Denial Date:	07/16/2015
Priority:	Standard	Application Received:	07/21/2015

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:

State(s) of Licensure: North Carolina

Certification(s)/Specialty: Family Practice

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The injured worker is a 50-year-old male who sustained a work related injury August 22, 2011. Electro diagnostic studies performed June 1, 2015; (report present in the medical record) revealed right C6 radiculitis, left C7 radiculitis, and moderate right and mild left carpal tunnel syndrome. According to the primary treating physician's progress report, dated July 8, 2015, the injured worker presented for evaluation regarding his neck, left shoulder, bilateral knees and low back pain. He has completed four of six acupuncture treatments for his neck and has noticed a decrease in pain. He continues with his home exercise program and his medications are helpful and well tolerated. Current medication included Celebrex, Omeprazole, and Nucynta. He rates his overall pain 6-7 out of 10 and is described as stabbing and aching in the neck, aching in the upper back, aching and burning in his shoulders, stabbing and aching in the lower back, burning and aching in the knees, stabbing and burning in his left foot, aching and burning in his left ankle, and aching and numbing in his wrists. He is unchanged since his last appointment. Physical examination revealed sensation is diminished into the left C6-7 dermatome, Spurling's test is negative but elicits pain, and Hoffman's negative bilaterally. There is mild swelling and effusion of the bilateral knees, joint line tenderness bilaterally; crepitus appreciated with flexion and extension, and McMurray's test is negative on the right and positive on the left. Impression is documented as shoulder pain' cervicalgia; pain of cervical facet joint; myalgia and myositis; lumbago; degeneration of lumbar intervertebral disc; thoracic or lumbosacral neuritis or radiculitis, unspecified; chronic pain syndrome. At issue, is a request for authorization for Nucynta.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

Nucynta 50mg, #90: Upheld

Claims Administrator guideline: The Claims Administrator did not base their decision on the MTUS. Decision based on Non-MTUS Citation Official Disability Guidelines (ODG-TWC) Treatment Integrated/Disability Duration Guidelines, Pain (Chronic) - Tapentadol (Nucynta).

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines opioids Page(s): 76-84.

Decision rationale: The California chronic pain medical treatment guidelines section on opioids states for ongoing management: On-Going Management. Actions Should Include: (a) Prescriptions from a single practitioner taken as directed, and all prescriptions from a single pharmacy. (b) The lowest possible dose should be prescribed to improve pain and function. (c) Office: Ongoing review and documentation of pain relief, functional status, appropriate medication use, and side effects. Pain assessment should include current pain; the least reported pain over the period since last assessment; average pain; intensity of pain after taking the opioid; how long it takes for pain relief; and how long pain relief lasts. Satisfactory response to treatment may be indicated by the patient's decreased pain, increased level of function, or improved quality of life. Information from family members or other caregivers should be considered in determining the patient's response to treatment. The 4 A's for Ongoing Monitoring: Four domains have been proposed as most relevant for ongoing monitoring of chronic pain patients on opioids: pain relief, side effects, physical and psychosocial functioning, and the occurrence of any potentially aberrant (or non-adherent) drug-related behaviors. These domains have been summarized as the "4 A's" (analgesia, activities of daily living, adverse side effects, and aberrant drug taking behaviors). The monitoring of these outcomes over time should affect therapeutic decisions and provide a framework for documentation of the clinical use of these controlled drugs. (Passik, 2000) (d) Home: To aid in pain and functioning assessment, the patient should be requested to keep a pain diary that includes entries such as pain triggers, and incidence of end-of-dose pain. It should be emphasized that using this diary will help in tailoring the opioid dose. This should not be a requirement for pain management. (e) Use of drug screening or inpatient treatment with issues of abuse, addiction, or poor pain control. (f) Documentation of misuse of medications (doctor- shopping, uncontrolled drug escalation, drug diversion). (g) Continuing review of overall situation with regard to non-opioid means of pain control. (h) Consideration of a consultation with a multidisciplinary pain clinic if doses of opioids are required beyond what is usually required for the condition or pain does not improve on opioids in 3 months. Consider a psych consult if there is evidence of depression, anxiety or irritability. Consider an addiction medicine consult if there is evidence of substance misuse. When to Continue Opioids (a) If the patient has returned to work. (b) If the patient has improved functioning and pain. (Washington, 2002) (Colorado, 2002) (Ontario, 2000) (VA/DoD, 2003) (Maddox-AAPM/APS, 1997) (Wisconsin, 2004) (Warfield, 2004) The long-term use of this medication class is not recommended per the California MTUS unless there documented evidence of benefit with measurable outcome measures and improvement in function. There is no documented significant decrease in objective pain measures such as VAS scores for significant periods of time. There are no objective measures of improvement of function. Therefore, not all criteria for the ongoing use of opioids have been met and the request is not medically necessary.

