

Case Number:	CM15-0144556		
Date Assigned:	08/05/2015	Date of Injury:	06/06/2003
Decision Date:	09/02/2015	UR Denial Date:	07/07/2015
Priority:	Standard	Application Received:	07/27/2015

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:

State(s) of Licensure: North Carolina

Certification(s)/Specialty: Family Practice

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The injured worker is a 66 year old male, who sustained an industrial injury on 6-06-2003. The mechanism of injury was not noted. The injured worker was diagnosed as having lumbar post-laminectomy syndrome and myofascial pain syndrome, low back. Treatment to date has included diagnostics, trigger point injections, and medications. Trigger point injections over the right and left low back and buttocks were administered on 4-21-2015. Medications at that time included Lyrica, Tramadol, and Lidoderm patches. Currently (6-22-2015), the injured worker complains of diffuse body pain, in addition to his low back pain radiating to his lower extremities, and pain in his neck radiating to his upper extremities. His pain level remained 7 out of 10 and was unchanged from previous visit. His diabetes was controlled and he requested trigger point injections. Exam of the thoracolumbar spine noted discrete tender trigger points over his low back and buttocks, with muscle twitch points, and decreased range of motion. Motor and sensation was decreased in the left L5-S1 distribution. Trigger point injections were administered over the right and left low back and buttocks. He remained on Lyrica, Tramadol, and Lidoderm patches, noting that the Lidoderm was not approved. A CURES report was documented as consistent and urine toxicology was not noted. He was documented to be at maximum medical improvement.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

Tramadol 50mg #120, plus 2 refills: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Opioids Page(s): 75, 78.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines opioids Page(s): 76-84.

Decision rationale: The California chronic pain medical treatment guidelines section on opioids states for ongoing management: On-Going Management. Actions Should Include: (a) Prescriptions from a single practitioner taken as directed, and all prescriptions from a single pharmacy.(b) The lowest possible dose should be prescribed to improve pain and function. (c) Office: Ongoing review and documentation of pain relief, functional status, appropriate medication use, and side effects. Pain assessment should include: current pain; the least reported pain over the period since last assessment; average pain; intensity of pain after taking the opioid; how long it takes for pain relief; and how long pain relief lasts. Satisfactory response to treatment may be indicated by the patient's decreased pain, increased level of function, or improved quality of life. Information from family members or other caregivers should be considered in determining the patient's response to treatment. The 4 A's for Ongoing Monitoring: Four domains have been proposed as most relevant for ongoing monitoring of chronic pain patients on opioids: pain relief, side effects, physical and psychosocial functioning, and the occurrence of any potentially aberrant (or non-adherent) drug-related behaviors. These domains have been summarized as the "4 A's" (analgesia, activities of daily living, adverse side effects, and aberrant drug taking behaviors). The monitoring of these outcomes over time should affect therapeutic decisions and provide a framework for documentation of the clinical use of these controlled drugs. (Passik, 2000) (d) Home: To aid in pain and functioning assessment, the patient should be requested to keep a pain diary that includes entries such as pain triggers, and incidence of end-of-dose pain. It should be emphasized that using this diary will help in tailoring the opioid dose. This should not be a requirement for pain management. (e) Use of drug screening or inpatient treatment with issues of abuse, addiction, or poor pain control. (f) Documentation of misuse of medications (doctor- shopping, uncontrolled drug escalation, drug diversion). (g) Continuing review of overall situation with regard to non-opioid means of pain control. (h) Consideration of a consultation with a multidisciplinary pain clinic if doses of opioids are required beyond what is usually required for the condition or pain does not improve on opioids in 3 months. Consider a psych consult if there is evidence of depression, anxiety or irritability. Consider an addiction medicine consult if there is evidence of substance misuse. When to Continue Opioids (a) If the patient has returned to work (b) If the patient has improved functioning and pain(Washington, 2002) (Colorado, 2002) (Ontario, 2000) (VA/DoD, 2003) (Maddox-AAPM/APS, 1997) (Wisconsin, 2004) (Warfield, 2004) The long-term use of this medication class is not recommended per the California MTUS unless there documented evidence of benefit with measurable outcome measures and improvement in function. There is no documented significant improvement in VAS scores for significant periods of time. There are no objective measurements of improvement in function. Therefore all criteria for the ongoing use of opioids have not been met and the request is not certified.

Retro Trigger point injections, lumbar spine, series of 4 DOS: 6.22.15: Overturned

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Page(s): 122.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines trigger point injections Page(s): 122.

Decision rationale: The California chronic pain medical treatment guidelines section on trigger point injections states: Trigger point injections Recommended only for myofascial pain syndrome as indicated below, with limited lasting value. Not recommended for radicular pain. Trigger point injections with an anesthetic such as bupivacaine are recommended for non-resolving trigger points, but the addition of a corticosteroid is not generally recommended. Not recommended for radicular pain. A trigger point is a discrete focal tenderness located in a palpable taut band of skeletal muscle, which produces a local twitch in response to stimulus to the band. Trigger points may be present in up to 33-50% of the adult population. Myofascial pain syndrome is a regional painful muscle condition with a direct relationship between a specific trigger point and its associated pain region. These injections may occasionally be necessary to maintain function in those with myofascial problems when myofascial trigger points are present on examination. Not recommended for typical back pain or neck pain. (Graff-Radford, 2004) (Nelemans-Cochrane, 2002) For fibromyalgia syndrome, trigger point injections have not been proven effective. (Goldenberg, 2004) Criteria for the use of Trigger point injections: Trigger point injections with a local anesthetic may be recommended for the treatment of chronic low back or neck pain with myofascial pain syndrome when all of the following criteria are met: (1) Documentation of circumscribed trigger points with evidence upon palpation of a twitch response as well as referred pain; (2) Symptoms have persisted for more than three months; (3) Medical management therapies such as ongoing stretching exercises, physical therapy, NSAIDs and muscle relaxants have failed to control pain; (4) Radiculopathy is not present (by exam, imaging, or neuro-testing); (5) Not more than 3-4 injections per session; (6) No repeat injections unless a greater than 50% pain relief is obtained for six weeks after an injection and there is documented evidence of functional improvement; (7) Frequency should not be at an interval less than two months; (8) Trigger point injections with any substance (e.g., saline or glucose) other than local anesthetic with or without steroid are not recommended. (Colorado, 2002) (BlueCross BlueShield, 2004) The provided clinical documentation and physical exam findings meets criteria as outlined above and therefore the request is certified.

Lyrica 50mg #120, plus 2 refills: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines lyrica Page(s): 19.

Decision rationale: The California chronic pain medical treatment guidelines section on Lyrica states: Pregabalin (Lyrica, no generic available) has been documented to be effective in treatment of diabetic neuropathy and post herpetic neuralgia, has FDA approval for both

indications, and is considered first-line treatment for both. This medication is designated as a Schedule V controlled substance because of its causal relationship with euphoria. (Blommel, 2007) This medication also has an anti-anxiety effect. Pregabalin is being considered by the FDA as treatment for generalized anxiety disorder and social anxiety disorder. In June 2007, the FDA announced the approval of pregabalin as the first approved treatment for fibromyalgia. (ICSI, 2007) (Tassone, 2007) (Knotkova, 2007) (Eisenberg, 2007) (Crofford, 2005) (Stacey, 2008) The patient does not have the diagnoses of diabetic neuropathy, fibromyalgia or post herpetic neuropathy. There is no documentation of failure of other first line agents for peripheral neuropathy. Therefore, guideline recommendations have not been met and the request is not certified.