

Case Number:	CM15-0145614		
Date Assigned:	08/06/2015	Date of Injury:	03/21/2010
Decision Date:	09/02/2015	UR Denial Date:	06/29/2015
Priority:	Standard	Application Received:	07/27/2015

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:

State(s) of Licensure: California

Certification(s)/Specialty: Physical Medicine & Rehabilitation

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The injured worker is a 37 year old male, who sustained an industrial injury on 03-21-2010. On provider visit dated 04-02-2015 the injured worker's examination revealed tenderness along the olecranon tip. Tinel's sign on the right elbow was noted. Reverse Phalen's test revealed numbness along the little finger on the left. Tenderness was noted over the medial epicondyle on the right was noted. On the left there was tenderness along the palmar ulnar carpal joint as well. The diagnoses have included lateral epicondylitis on the right, cubital tunnel syndrome on the right status post decompression and epicondylar release, ulnar impaction along the wrist and on the left and due to chronic pain the injured worker was noted to have an element of depression. Treatment to date has included medication, left wrist brace, elbow splint, elbow sleeve and TENS unit. The provider requested Protonix and Tramadol ER.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

Protonix 20mg #60: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines NSAIDs, GI symptoms & cardiovascular risk Page(s): 68.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines NSAIDs, GI Symptoms and Cardiovascular risk, Pages 68-69.

Decision rationale: Proton pump inhibitor (PPI) medication is for treatment of the problems associated with active gastric ulcers, erosive esophagitis, Barrett's esophagitis, or in patients with pathologic hypersecretion diseases. Although preventive treatment is effective for the mentioned diagnosis, studies suggest; however, nearly half of PPI prescriptions are used for unapproved or no indications. Per MTUS Chronic Pain Treatment Guidelines, the patient does not meet criteria for Omeprazole (Prilosec) namely reserved for patients with history of prior GI bleeding, the elderly (over 65 years), diabetics, and chronic cigarette smokers. Long term use of PPIs have potential increased risks of B12 deficiency; iron deficiency; hypomagnesemia; susceptibility to pneumonia, enteric infections, fractures, hypergastrinemia and cancer, and cardiovascular effects of myocardial infarction (MI). In the elderly, studies have demonstrated increased risk for Clostridium difficile infection, bone loss, and fractures from long-term use of PPIs. Given treatment criteria outweighing risk factors, if a PPI is to be used, omeprazole (Prilosec), lansoprazole (Prevacid), and esomeprazole (Nexium) are to be considered over second-line therapy of other PPIs such as pantoprazole (Protonix), dexlansoprazole (Dexilant), and rabeprazole (Aciphex). Submitted reports have not described or provided any GI diagnosis that meets the criteria to indicate medical treatment. Review of the records show no documentation of any specific history, identified symptoms, or confirmed GI diagnosis to warrant this medication. The Protonix 20mg #60 is not medically necessary and appropriate.

Tramadol ER 150mg #30: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Tramadol Opioids Page(s): 93-94,78-80 and 124.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Opioids, page(s) 74-96.

Decision rationale: The MTUS Guidelines cite opioid use in the setting of chronic, non-malignant, or neuropathic pain is controversial. Patients on opioids should be routinely monitored for signs of impairment and use of opioids in patients with chronic pain should be reserved for those with improved functional outcomes attributable to their use, in the context of an overall approach to pain management that also includes non-opioid analgesics, adjuvant therapies, psychological support, and active treatments (e.g., exercise). Submitted documents show no evidence that the treating physician is prescribing opioids in accordance to change in pain relief, functional goals with demonstrated improvement in daily activities, decreased in medical utilization or change in functional status. There is no evidence presented of random drug testing results or utilization of pain contract to adequately monitor for narcotic safety, efficacy, and compliance. The MTUS provides requirements of the treating physician to assess and document for functional improvement with treatment intervention and maintenance of function that would otherwise deteriorate if not supported. From the submitted reports, there is no demonstrated evidence of specific functional benefit derived from the continuing use of opioids with persistent severe pain for this chronic injury without acute flare, new injury, or progressive deterioration. The Tramadol ER 150mg #30 is not medically necessary and appropriate.