

Case Number:	CM15-0145894		
Date Assigned:	08/07/2015	Date of Injury:	10/06/2010
Decision Date:	09/04/2015	UR Denial Date:	07/06/2015
Priority:	Standard	Application Received:	07/28/2015

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:
 State(s) of Licensure: New Jersey, Alabama, California
 Certification(s)/Specialty: Neurology, Neuromuscular Medicine

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The injured worker is a 51-year-old male with an industrial injury dated 10-06-2010. The injured worker's diagnoses include bilateral trigger thumbs and status post bilateral trigger thumb release. Treatment consisted of diagnostic studies, prescribed medications, and periodic follow up visits. In a progress note dated 06-23-2015, the injured worker presented for treatment of bilateral hand injury. The injured worker rated pain a 6 out of 10. The injured worker reported improvement with swimming, stretching, hot weather and topical gel. The injured worker also reported that the pain increases with prolonged repetitive use, clicking activities and cold weather. Objective findings revealed tenderness to palpitation over the flexor and extensor tendon left thumb and slightly decreased motor strength in the left hand. The treating physician reported that the injured worker was experiencing a flare-up in the tendinitis of his left thumb with a treatment plan consisting of topical gel and physical therapy. The treating physician requested services for physical therapy evaluation x1 and treatment 2x4 for bilateral thumbs, now under review.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

PT Eval x1 and Treat 2x4 Bilateral Thumbs: Upheld

Claims Administrator guideline: The Claims Administrator did not base their decision on the MTUS. Decision based on Non-MTUS Citation Official Disability Guidelines (ODG), Forearm, Hand and Wrist chapter.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Physical Medicine Page(s): 98.

Decision rationale: According to MTUS guidelines, Physical Medicine is "Recommended as indicated below. Passive therapy (those treatment modalities that do not require energy expenditure on the part of the patient) can provide short-term relief during the early phases of pain treatment and are directed at controlling symptoms such as pain, inflammation and swelling and to improve the rate of healing soft tissue injuries. They can be used sparingly with active therapies to help control swelling, pain and inflammation during the rehabilitation process. Active therapy is based on the philosophy that therapeutic exercise and/or activity are beneficial for restoring flexibility, strength, endurance, function, range of motion, and can alleviate discomfort. Active therapy requires an internal effort by the individual to complete a specific exercise or task. This form of therapy may require supervision from a therapist or medical provider such as verbal, visual and/or tactile instruction(s). Patients are instructed and expected to continue active therapies at home as an extension of the treatment process in order to maintain improvement levels. Home exercise can include exercise with or without mechanical assistance or resistance and functional activities with assistive devices. (Colorado, 2002) (Airaksinen, 2006) Patient-specific hand therapy is very important in reducing swelling, decreasing pain, and improving range of motion in CRPS. (Li, 2005) The use of active treatment modalities (e.g., exercise, education, activity modification) instead of passive treatments is associated with substantially better clinical outcomes. In a large case series of patients with low back pain treated by physical therapists, those adhering to guidelines for active rather than passive treatments incurred fewer treatment visits, cost less, and had less pain and less disability. The overall success rates were 64.7% among those adhering to the active treatment recommendations versus 36.5% for passive treatment. (Fritz, 2007)" In this case, the patient underwent a right trigger thumb release on July 23, 2014 and a left trigger thumb release on April 23, 2014. The patient has had 8 post-op physical therapy sessions without clear documentation of efficacy. There is no documentation that the patient cannot perform home exercise program. Therefore, the request for 1 physical therapy evaluation and 8 physical therapy sessions for bilateral thumbs is not medically necessary.