

Case Number:	CM15-0146548		
Date Assigned:	08/07/2015	Date of Injury:	05/25/2005
Decision Date:	09/03/2015	UR Denial Date:	07/15/2015
Priority:	Standard	Application Received:	07/28/2015

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:
 State(s) of Licensure: North Carolina
 Certification(s)/Specialty: Family Practice

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The injured worker is a 63 year old male, who sustained an industrial injury on 5-25-2005. He reported an initial fall with injury requiring surgery to the right foot with cumulative traumatic injury to the low back, left hip, and bilateral upper extremities. Diagnoses include partial rotator cuff tear, pain in shoulder joint, hip joint replacement, chronic pain due to trauma and depression and anxiety. He has had multiple surgeries, two left hip surgeries, two shoulder repair, bilateral carpal tunnel repair, bilateral tendonitis repair, and trochanteric bursectomy of the greater trochanteric region. Treatments to date include medication therapy, activity modification, physical therapy, and psychotherapy. Currently, he complained of increased depression, stress, and short temper. He reported sleeping 4 to 5 hours at night. On 7-1-15, the physical examination documented fair concentration. The plan of care included prescription for Klonopin 0.05mg #60 and Restoril 30mg #30.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

Klonopin 0.5mg #60: Upheld

Claims Administrator guideline: The Claims Administrator did not cite any medical evidence for its decision.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines benzodiazepines Page(s): 24.

Decision rationale: The California chronic pain medical treatment guidelines section on benzodiazepines states: Benzodiazepines Not recommended for long-term use because long-term efficacy is unproven and there is a risk of dependence. Most guidelines limit use to 4 weeks. Their range of action includes sedative/hypnotic, anxiolytic, anticonvulsant, and muscle relaxant. Chronic benzodiazepines are the treatment of choice in very few conditions. Tolerance to hypnotic effects develops rapidly. Tolerance to anxiolytic effects occurs within months and long-term use may actually increase anxiety. A more appropriate treatment for anxiety disorder is an antidepressant. Tolerance to anticonvulsant and muscle relaxant effects occurs within weeks. (Baillargeon, 2003) (Ashton, 2005) The chronic long-term use of this class of medication is recommended in very few conditions per the California MTUS. There is no evidence however of all failure of first line agent for the treatment of anxiety or Insomnia in the provided documentation. For this reason the request is not medically necessary.

Restoril 30mg #30: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Benzodiazepines Page(s): 24.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines benzodiazepines Page(s): 24.

Decision rationale: The California chronic pain medical treatment guidelines section on benzodiazepines states: Benzodiazepines Not recommended for long-term use because long-term efficacy is unproven and there is a risk of dependence. Most guidelines limit use to 4 weeks. Their range of action includes sedative/hypnotic, anxiolytic, anticonvulsant, and muscle relaxant. Chronic benzodiazepines are the treatment of choice in very few conditions. Tolerance to hypnotic effects develops rapidly. Tolerance to anxiolytic effects occurs within months and long-term use may actually increase anxiety. A more appropriate treatment for anxiety disorder is an antidepressant. Tolerance to anticonvulsant and muscle relaxant effects occurs within weeks. (Baillargeon, 2003) (Ashton, 2005) The chronic long-term use of this class of medication is recommended in very few conditions per the California MTUS. There is no evidence however of all failure of first line agent for the treatment of anxiety or Insomnia in the provided documentation. For this reason the request is not medically necessary.