

Case Number:	CM15-0146655		
Date Assigned:	08/07/2015	Date of Injury:	09/17/2012
Decision Date:	09/04/2015	UR Denial Date:	07/13/2015
Priority:	Standard	Application Received:	07/28/2015

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:

State(s) of Licensure: North Carolina

Certification(s)/Specialty: Family Practice

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The injured worker is a 57-year-old female, who sustained an industrial injury on 09-17-2012 secondary to lifting a person resulting in low back pain. On provider visit dated 06-29-2015 the injured worker has reported ongoing low back pain and epigastric pain with nausea. Objective findings were noted as ongoing tenderness to the lumbar paraspinal muscles and one point tenderness in the epigastric area. The diagnoses have included chronic low back pain. Treatment to date has included medication currently listed as Norco, Prilosec and Colace. The provider requested Colace and Prilosec.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

Colace QID #120 refills 2: Overturned

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Criteria for use of opioids. Decision based on Non-MTUS Citation Official Disability Guidelines, Opioid induced constipation treatment.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines opioids
Page(s): 77.

Decision rationale: The California chronic pain medical treatment guidelines section on opioid therapy states: (a) Intermittent pain: Start with a short-acting opioid trying one medication at a time. (b) Continuous pain: extended-release opioids are recommended. Patients on this modality may require a dose of "rescue" opioids. The need for extra opioid can be a guide to determine the sustained release dose required. (c) Only change 1 drug at a time. (d) Prophylactic treatment of constipation should be initiated. The patient is currently on opioid therapy. The use of constipation measures is advised per the California MTUS. The requested medication is used in the treatment of constipation. Therefore, the request is medically necessary.

Prilosec 20mg QD #30 refills 2: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines NSAIDs, GI & Cardiovascular Risk.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines NSAID Page(s): 68.

Decision rationale: The California chronic pain medical treatment guidelines section on NSAID therapy and proton pump inhibitors (PPI) states: Recommend with precautions as indicated below. Clinicians should weight the indications for NSAIDs against both GI and cardiovascular risk factors. Determine if the patient is at risk for gastrointestinal events: (1) age > 65 years; (2) history of peptic ulcer, GI bleeding or perforation; (3) concurrent use of ASA, corticosteroids, and/or an anticoagulant; or (4) high dose/multiple NSAID (e.g., NSAID + low-dose ASA). Recent studies tend to show that H. Pylori does not act synergistically with NSAIDS to develop gastro duodenal lesions. Recommendations: Patients with no risk factor and no cardiovascular disease: Non-selective NSAIDs OK (e.g, ibuprofen, naproxen, etc.). Patients at intermediate risk for gastrointestinal events and no cardiovascular disease: (1) A non-selective NSAID with either a PPI (Proton Pump Inhibitor, for example, 20 mg omeprazole daily) or misoprostol (200 ug four times daily) or (2) a Cox-2 selective agent. Long-term PPI use (> 1 year) has been shown to increase the risk of hip fracture (adjusted odds ratio 1.44). Patients at high risk for gastrointestinal events with no cardiovascular disease: A Cox-2 selective agent plus a PPI if absolutely necessary. There is no documentation provided that places this patient at intermediate or high risk that would justify the use of a PPI. There is no mention of current gastrointestinal or cardiovascular disease. For these reasons, the criteria set forth above per the California MTUS for the use of this medication has not been met. Therefore, the request is not medically necessary.