

<b>Case Number:</b>	CM15-0163426		
<b>Date Assigned:</b>	08/31/2015	<b>Date of Injury:</b>	11/29/2012
<b>Decision Date:</b>	10/14/2015	<b>UR Denial Date:</b>	07/18/2015
<b>Priority:</b>	Standard	<b>Application Received:</b>	08/20/2015

### HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:

State(s) of Licensure: California, District of Columbia, Maryland

Certification(s)/Specialty: Anesthesiology, Pain Management

### CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

This 41 year old man sustained an industrial injury on 11-19-2012 while pulling the rubber back from a mold. He felt immediate pain I his hand and it swelled up. Evaluations include left wrist x-rays dated 3-25-2015 and left elbow MRI dated 1-11-2013. Diagnoses include left wrist severe ankyloses and carpal tunnel dissection. Treatment has included oral medications and physical therapy. Physician notes dated 7-13-2015 show complaints of left wrist and hand pain rated 9 out of 10 with numbness, weakness, stiffness, popping, grinding, and swelling. Recommendations include hand specialist consultation, pain management consultation, rheumatology consultation, psychological evaluation, left wrist CT scan, left wrist brace, Naproxen, Pantoprazole, Norco, and follow up in six weeks.

### IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

**Flexeril 7.5 mg, unknown quantity:** Upheld

**Claims Administrator guideline:** Decision based on MTUS Chronic Pain Medical Treatment 2009.

**MAXIMUS guideline:** Decision based on MTUS Chronic Pain Medical Treatment 2009, Section(s): Muscle relaxants (for pain).

**Decision rationale:** With regard to muscle relaxants, the MTUS CPMTG states: "Recommend non-sedating muscle relaxants with caution as a second-line option for short-term treatment of acute exacerbations in patients with chronic LBP. (Chou, 2007) (Mens, 2005) (Van Tulder, 1998) (van Tulder, 2003) (van Tulder, 2006) (Schnitzer, 2004) (See, 2008) Muscle relaxants may be effective in reducing pain and muscle tension, and increasing mobility. However, in most LBP cases, they show no benefit beyond NSAIDs in pain and overall improvement." Regarding Cyclobenzaprine: "Recommended for a short course of therapy. Limited, mixed-evidence does not allow for a recommendation for chronic use. Cyclobenzaprine is a skeletal muscle relaxant and a central nervous system depressant with similar effects to tricyclic antidepressants (e.g. amitriptyline). Cyclobenzaprine is more effective than placebo in the management of back pain, although the effect is modest and comes at the price of adverse effects." Per p41 of the MTUS guidelines the effect is greatest in the first 4 days of treatment, suggesting that shorter courses may be better. Treatment is recommended for the treatment of acute spasm limited to a maximum of 2-3 weeks. UDS that evaluate for cyclobenzaprine can provide additional data on whether the injured worker is compliant, however in this case there is no UDS testing for cyclobenzaprine. The documentation submitted for review indicates that the injured worker has been using this medication since at least 6-2015. There is no documentation of the patients' specific functional level or percent improvement with treatment with cyclobenzaprine. As it is recommended only for short-term use, medical necessity cannot be affirmed. Furthermore, the request does not specify quantity information. Therefore, the request is not medically necessary.

**Norco 10/325 mg, unknown quantity:** Upheld

**Claims Administrator guideline:** Decision based on MTUS Chronic Pain Medical Treatment 2009.

**MAXIMUS guideline:** Decision based on MTUS Chronic Pain Medical Treatment 2009, Section(s): Opioids, criteria for use.

**Decision rationale:** Per MTUS Chronic Pain Medical Treatment Guidelines p78 regarding on-going management of opioids "Four domains have been proposed as most relevant for ongoing monitoring of chronic pain patients on opioids: Pain relief, side effects, physical and psychosocial functioning, and the occurrence of any potentially aberrant (or non-adherent) drug related behaviors. These domains have been summarized as the "4 A's" (Analgesia, activities of daily living, adverse side effects, and any aberrant drug-taking behaviors). The monitoring of these outcomes over time should affect therapeutic decisions and provide a framework for documentation of the clinical use of these controlled drugs." Review of the available medical records reveals no documentation to support the medical necessity of Norco nor any documentation addressing the '4 A's' domains, which is a recommended practice for the on-going management of opioids. Specifically, the notes do not appropriately review and document pain relief, functional status improvement, appropriate medication use, or side effects. The MTUS considers this list of criteria for initiation and continuation of opioids in the context of efficacy required to substantiate medical necessity, and they do not appear to have been addressed by the treating physician in the documentation available for review. Efforts to rule out aberrant behavior (e.g. CURES report, UDS, opiate agreement) are necessary to assure safe usage and establish medical necessity. There is no documentation comprehensively addressing this concern in the records available for my review. As MTUS recommends to discontinue opioids if there is no overall improvement in function, medical necessity cannot be affirmed. Furthermore, the request does not specify quantity information. Therefore, the request is not medically necessary.

**Ultram ER 150 mg, unknown quantity:** Upheld

**Claims Administrator guideline:** Decision based on MTUS Chronic Pain Medical Treatment 2009.

**MAXIMUS guideline:** Decision based on MTUS Chronic Pain Medical Treatment 2009, Section(s): Opioids, criteria for use.

**Decision rationale:** Per MTUS Chronic Pain Medical Treatment Guidelines p78 regarding ongoing management of opioids "Four domains have been proposed as most relevant for ongoing monitoring of chronic pain patients on opioids: Pain relief, side effects, physical and psychosocial functioning, and the occurrence of any potentially aberrant (or non-adherent) drug related behaviors. These domains have been summarized as the "4 A's" (Analgesia, activities of daily living, adverse side effects, and any aberrant drug-taking behaviors). The monitoring of these outcomes over time should affect therapeutic decisions and provide a framework for documentation of the clinical use of these controlled drugs." Review of the available medical records reveals no documentation to support the medical necessity of Ultram nor any documentation addressing the '4 A's' domains, which is a recommended practice for the ongoing management of opioids. Specifically, the notes do not appropriately review and document pain relief, functional status improvement, appropriate medication use, or side effects. The MTUS considers this list of criteria for initiation and continuation of opioids in the context of efficacy required to substantiate medical necessity, and they do not appear to have been addressed by the treating physician in the documentation available for review. Furthermore, efforts to rule out aberrant behavior (e.g. CURES report, UDS, opiate agreement) are necessary to assure safe usage and establish medical necessity. There is no documentation comprehensively addressing this concern in the records available for my review. As MTUS recommends to discontinue opioids if there is no overall improvement in function, medical necessity cannot be affirmed. Furthermore, the request does not specify quantity information. Therefore, the request is not medically necessary.

**Voltaren XR 100 mg, unknown quantity:** Upheld

**Claims Administrator guideline:** Decision based on MTUS Chronic Pain Medical Treatment 2009.

**MAXIMUS guideline:** Decision based on MTUS Chronic Pain Medical Treatment 2009, Section(s): NSAIDs (non-steroidal anti-inflammatory drugs).

**Decision rationale:** With regard to the use of NSAIDs for chronic low back pain, the MTUS CPMTG states "Recommended as an option for short-term symptomatic relief. A Cochrane review of the literature on drug relief for low back pain (LBP) suggested that NSAIDs were no more effective than other drugs such as acetaminophen, narcotic analgesics, and muscle relaxants. The review also found that NSAIDs had more adverse effects than placebo and acetaminophen but fewer effects than muscle relaxants and narcotic analgesics. In addition, evidence from the review suggested that no one NSAID, including COX-2 inhibitors, was clearly more effective than another." "Low back pain (chronic): Both acetaminophen and NSAIDs have been recommended as first line therapy for low back pain. There is insufficient evidence to recommend one medication over the other. Selection should be made on a case-by-case basis based on weighing efficacy vs. side effect profile." The documentation submitted for review indicates that the injured worker has been using NSAIDs since at least 6-2015. As the request does not specify quantity information, medical necessity cannot be affirmed. Therefore, the request is not medically necessary.

**Prilosec 20 mg, unknown quantity:** Upheld

**Claims Administrator guideline:** Decision based on MTUS Chronic Pain Medical Treatment 2009.

**MAXIMUS guideline:** Decision based on MTUS Chronic Pain Medical Treatment 2009, Section(s): NSAIDs, GI symptoms & cardiovascular risk.

**Decision rationale:** In the treatment of dyspepsia secondary to NSAID therapy, the MTUS recommends stopping the NSAID, switching to a different NSAID, or considering the use of an H2-receptor antagonist or a PPI. The MTUS Chronic Pain Medical Treatment Guidelines recommend the use of proton pump inhibitors in conjunction with NSAIDs in situations in which the patient is at risk for gastrointestinal events including: (1) age > 65 years; (2) history of peptic ulcer, GI bleeding or perforation; (3) concurrent use of ASA, corticosteroids, and-or an anticoagulant; or (4) high dose-multiple NSAID (e.g., NSAID + low-dose ASA). CPMTG guidelines further specify: "Recommendations: Patients with no risk factor and no cardiovascular disease: Non-selective NSAIDs OK (e.g., ibuprofen, naproxen, etc.). Patients at intermediate risk for gastrointestinal events and no cardiovascular disease: (1) A non-selective NSAID with either a PPI (Proton Pump Inhibitor, for example, 20 mg omeprazole daily) or misoprostol (200g four times daily) or (2) a Cox-2 selective agent. Long-term PPI use (> 1 year) has been shown to increase the risk of hip fracture (adjusted odds ratio 1.44). Patients at high risk for gastrointestinal events with no cardiovascular disease: A Cox-2 selective agent plus a PPI if absolutely necessary. Patients at high risk of gastrointestinal events with cardiovascular disease: If GI risk is high the suggestion is for a low-dose Cox-2 plus low dose Aspirin (for cardioprotection) and a PPI. If cardiovascular risk is greater than GI risk, the suggestion is naproxyn plus low-dose aspirin plus a PPI. (Laine, 2006) (Scholmerich, 2006) (Nielsen, 2006) (Chan, 2004) (Gold, 2007) (Laine, 2007)." As continued NSAID therapy is not medically necessary and there is no documentation of peptic ulcer, GI bleeding or perforation, or cardiovascular disease in the records available for my review, the injured worker's risk for gastrointestinal events is low, as such, medical necessity cannot be affirmed. Furthermore, the request does not specify quantity information. Therefore, the request is not medically necessary.

