MAXIMUS Federal Services

Case Number:	CM15-0165341		
Date Assigned:	09/02/2015	Date of Injury:	02/04/2015
Decision Date:	10/06/2015	UR Denial Date:	07/21/2015
Priority:	Standard	Application Received:	08/24/2015

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:
State(s) of Licensure: North Carolina
Certification(s)/Specialty: Family Practice

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The injured worker is a 45-year-old female with an industrial injury dated 02-04-2015. His diagnoses included lateral epicondylitis right elbow and radial tunnel syndrome right proximal forearm. Comorbid conditions included hypothyroidism, hyperlipidemia and hypertension. Prior treatment included physical therapy, elbow cuff, Flector patches and occupational therapy. She presents on 07-10-2015 with complaints of pain over the right proximal radial forearm in the area of the radial tunnel. Physical findings included maximal tenderness to palpation over the mid, right and proximal dorsal forearm. She was no longer tender over the lateral epicondyle and mid dorsal right wrist. She still had pain in the proximal right forearm with supination against resistance. The provider documents improved pain level from 10 out of 10 to 6 out of 10 for at least 24 hours following each session. She had also been able to wash dishes and vacuum. The provider documented in light of the positive response 8 more physical therapy visits are being requested to decrease her pain and increase her ability to function and strengthen her right arm enough to allow return to work. The treatment request is for 8 additional visits of physical therapy (frequency and duration unspecified) for the right elbow.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

8 additional visits of Physical therapy (frequency and duration unspecified) for the right elbow: Upheld

Claims Administrator guideline: Decision based on MTUS ACOEM Chapter 10 Elbow Disorders (Revised 2007), Chapter 11 Forearm, Wrist, and Hand Complaints, Chronic Pain Treatment Guidelines.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines physical medicine Page(s): 98-99.

Decision rationale: The California chronic pain medical treatment guidelines section on physical medicine states: Recommended as indicated below. Passive therapy (those treatment modalities that do not require energy expenditure on the part of the patient) can provide short-term relief during the early phases of pain treatment and are directed at controlling symptoms such as pain, inflammation and swelling and to improve the rate of healing soft tissue injuries. They can be used sparingly with active therapies to help control swelling, pain and inflammation during the rehabilitation process. Active therapy is based on the philosophy that therapeutic exercise and/or activity are beneficial for restoring flexibility, strength, endurance, function, range of motion, and can alleviate discomfort. Active therapy requires an internal effort by the individual to complete a specific exercise or task. This form of therapy may require supervision from a therapist or medical provider such as verbal, visual and/or tactile instruction(s). Patients are instructed and expected to continue active therapies at home as an extension of the treatment process in order to maintain improvement levels. Home exercise can include exercise with or without mechanical assistance or resistance and functional activities with assistive devices. (Colorado, 2002) (Airaksinen, 2006) Patient-specific hand therapy is very important in reducing swelling, decreasing pain, and improving range of motion in CRPS. (Li, 2005) The use of active treatment modalities (e.g., exercise, education, activity modification) instead of passive treatments is associated with substantially better clinical outcomes. In a large case series of patients with low back pain treated by physical therapists, those adhering to guidelines for active rather than passive treatments incurred fewer treatment visits, cost less, and had less pain and less disability. The overall success rates were 64.7% among those adhering to the active treatment recommendations versus 36.5% for passive treatment. (Fritz, 2007) Physical Medicine Guidelines Allow for fading of treatment frequency (from up to 3 visits per week to 1 or less), plus active self-directed home Physical Medicine. Myalgia and myositis, unspecified (ICD9 729.1): 9-10 visits over 8 weeks; Neuralgia, neuritis, and radiculitis, unspecified (ICD9 729.2) 8-10 visits over 4 weeks; Reflex sympathetic dystrophy (CRPS) (ICD9 337.2): 24 visits over 16 weeks. The goal of physical therapy is graduation to home therapy after a certain amount of recommended sessions. The patient has already completed physical therapy. The request is in excess of these recommendations per the California MTUS. There is no objective reason why the patient would not be moved to home therapy after completing the recommended amount of supervised sessions in the provided clinical documentation. Therefore, the request is not medically necessary.