

Case Number:	CM15-0165557		
Date Assigned:	09/03/2015	Date of Injury:	03/17/2015
Decision Date:	10/06/2015	UR Denial Date:	07/28/2015
Priority:	Standard	Application Received:	08/24/2015

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:
 State(s) of Licensure: North Carolina
 Certification(s)/Specialty: Family Practice

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The injured worker is a 63-year-old female, who sustained an industrial injury on 3-17-15. She has reported initial complaints of twisting the left ankle with injury striking the left side back and left hip. The diagnoses have included lumbar degenerative disc disease (DDD), lumbar sprain and strain, left hip contusion, left hip strain and sprain, left ankle sprain and strain, and left ankle rule out internal derangement. Treatment to date has included medications, diagnostics, physical therapy, work restrictions, and other modalities. Currently, as per the physician progress note dated 7-10-15, the injured worker complains of left ankle and foot pain that is on and off and associated with tingling. The diagnostic testing that was performed included Magnetic Resonance Imaging (MRI) of the left foot and ankle dated 6-6-15 that reveals mild strain of the ligaments and small joint effusion. There was also an X-ray of the left ankle dated 4-2-15. The objective findings-physical exam of the left foot and ankle reveals tenderness to palpation over the anterolateral ankle with swelling noted. There is acupuncture sessions noted. The physician requested treatments included Physical therapy twice a week for six weeks and Acupuncture once a week for six weeks to the left ankle and foot to decrease pain and increase range of motion.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

Physical therapy twice a week for six weeks: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Physical Medicine.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines physical medicine Page(s): 98-99.

Decision rationale: The California chronic pain medical treatment guidelines section on physical medicine states: Recommended as indicated below. Passive therapy (those treatment modalities that do not require energy expenditure on the part of the patient) can provide short-term relief during the early phases of pain treatment and are directed at controlling symptoms such as pain, inflammation and swelling and to improve the rate of healing soft tissue injuries. They can be used sparingly with active therapies to help control swelling, pain and inflammation during the rehabilitation process. Active therapy is based on the philosophy that therapeutic exercise and/or activity are beneficial for restoring flexibility, strength, endurance, function, range of motion, and can alleviate discomfort. Active therapy requires an internal effort by the individual to complete a specific exercise or task. This form of therapy may require supervision from a therapist or medical provider such as verbal, visual and/or tactile instruction(s). Patients are instructed and expected to continue active therapies at home as an extension of the treatment process in order to maintain improvement levels. Home exercise can include exercise with or without mechanical assistance or resistance and functional activities with assistive devices. (Colorado, 2002) (Airaksinen, 2006) Patient-specific hand therapy is very important in reducing swelling, decreasing pain, and improving range of motion in CRPS. (Li, 2005) The use of active treatment modalities (e.g., exercise, education, activity modification) instead of passive treatments is associated with substantially better clinical outcomes. In a large case series of patients with low back pain treated by physical therapists, those adhering to guidelines for active rather than passive treatments incurred fewer treatment visits, cost less, and had less pain and less disability. The overall success rates were 64.7% among those adhering to the active treatment recommendations versus 36.5% for passive treatment. (Fritz, 2007) Physical Medicine Guidelines. Allow for fading of treatment frequency (from up to 3 visits per week to 1 or less), plus active self-directed home Physical Medicine. Myalgia and myositis, unspecified (ICD9 729. 1): 9-10 visits over 8 weeks. Neuralgia, neuritis, and radiculitis, unspecified (ICD9 729. 2) 8-10 visits over 4 weeks. Reflex sympathetic dystrophy (CRPS) (ICD9 337. 2): 24 visits over 16 weeks. The goal of physical therapy is graduation to home therapy after a certain amount of recommended sessions. The patient has already completed physical therapy. The request is in excess of these recommendations per the California MTUS. There is no objective reason why the patient would not be moved to home therapy after completing the recommended amount of supervised sessions. In the provided clinical documentation. Therefore, the request is not medically necessary.

Acupuncture once a week for six weeks: Overturned

Claims Administrator guideline: Decision based on MTUS Acupuncture Treatment Guidelines.

MAXIMUS guideline: Decision based on MTUS Acupuncture Treatment Guidelines.

Decision rationale: The California chronic pain medical treatment guidelines section on acupuncture states: 1) "Acupuncture" is used as an option when pain medication is reduced or not tolerated, it may be used as an adjunct to physical rehabilitation and/or surgical intervention to hasten functional recovery. It is the insertion and removal of filiform needles to stimulate acupoints (acupuncture points). Needles may be inserted, manipulated, and retained for a period of time. Acupuncture can be used to reduce pain, reduce inflammation, increase blood flow, increase range of motion, decrease the side effect of medication-induced nausea, promote relaxation in an anxious patient, and reduce muscle spasm. Frequency and duration of acupuncture with electrical stimulation may be performed as follows: 1. Time to produce functional improvement 3-6 treatments. 2. Frequency: 1-3 times per week. 3. Optimum duration is 1-2 months. 4. Treatments may be extended if functional improvement is documented. The request for acupuncture is for a total of 6 sessions. This is not in excess of the recommendations. The patient must demonstrate functional improvement in 3-6 treatments for more sessions to be certified. Therefore, the request is not in excess of the recommended initial treatment sessions and is medically necessary.