

Case Number:	CM15-0165967		
Date Assigned:	09/03/2015	Date of Injury:	05/10/2004
Decision Date:	10/06/2015	UR Denial Date:	08/14/2015
Priority:	Standard	Application	08/24/2015

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:

State(s) of Licensure: North Carolina

Certification(s)/Specialty: Family Practice

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The injured worker (IW) is a 55-year-old male who sustained an industrial injury on 05-10-2004. The mechanism of the injury is not found in the records reviewed. The injured worker was diagnosed as having osteoarthritis of the knee, status post right knee partial meniscectomy 09/2004, and lumbar spine 3 mm disc bulge. Treatment to date has included surgery, and medications. Currently, the injured worker complains of low back pain rated a 4 on a scale of 01- 10 described as an ache, and bilateral hip pain rated a 1-2 on a scale of 0-10. Since bilateral hip surgeries, his pain has decreased significantly. He also had pain in the bilateral knees; right greater than left rated a 7-8 on a scale of 0-10. The knee pain was worsening with the right knee feeling like it was hyper extending and giving way. His pain is accompanied by numbness and tingling and is aggravated by prolonged standing, walking, walking on uneven surfaces, lifting heavy objects, and cold weather. The worker s treatment includes oral medications and topical analgesic creams. He states his pain does diminish with medications and analgesic topical creams. On exam, he has bilateral paraspinal tenderness L5-S1 and mild tenderness bilaterally at the S1. He has good range of motion in the lumbar spine. He has nonspecific tenderness at both knees and tenderness to palpation of the bilateral thighs. The treatment plan included medications, x-rays and MRI of the right knee, and refills on medications with consultation of a home exercise program. A request for authorization was submitted for Norco 10/325mg #120 and a consultation for home exercise program. A utilization review decision (08-14-2015) approved the consultation for a home exercise program, and declined the Norco request due to lack of documentation of the "4 A's" (Analgesia, Activities of daily living, Adverse side effects and Adverse behavior) in response to medication.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

Norco 10/325mg #120: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines opioids Page(s): 78.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines opioids Page(s): 76-84.

Decision rationale: The California chronic pain medical treatment guidelines section on opioids states for ongoing management: On-Going Management. Actions Should Include: (a) Prescriptions from a single practitioner taken as directed, and all prescriptions from a single pharmacy. (b) The lowest possible dose should be prescribed to improve pain and function. (c) Office: Ongoing review and documentation of pain relief, functional status, appropriate medication use, and side effects. Pain assessment should include: current pain; the least reported pain over the period since last assessment; average pain; intensity of pain after taking the opioid; how long it takes for pain relief; and how long pain relief lasts. Satisfactory response to treatment may be indicated by the patient's decreased pain, increased level of function, or improved quality of life. Information from family members or other caregivers should be considered in determining the patient's response to treatment. The 4 A's for Ongoing Monitoring: Four domains have been proposed as most relevant for ongoing monitoring of chronic pain patients on opioids: pain relief, side effects, physical and psychosocial functioning, and the occurrence of any potentially aberrant (or non-adherent) drug-related behaviors. These domains have been summarized as the "4 A's" (analgesia, activities of daily living, adverse side effects, and aberrant drug taking behaviors). The monitoring of these outcomes over time should affect therapeutic decisions and provide a framework for documentation of the clinical use of these controlled drugs. (Passik, 2000) (d) Home: To aid in pain and functioning assessment, the patient should be requested to keep a pain diary that includes entries such as pain triggers, and incidence of end-of-dose pain. It should be emphasized that using this diary will help in tailoring the opioid dose. This should not be a requirement for pain management. (e) Use of drug screening or inpatient treatment with issues of abuse, addiction, or poor pain control. (f) Documentation of misuse of medications (doctor- shopping, uncontrolled drug escalation, drug diversion). (g) Continuing review of overall situation with regard to non-opioid means of pain control. (h) Consideration of a consultation with a multidisciplinary pain clinic if doses of opioids are required beyond what is usually required for the condition or pain does not improve on opioids in 3 months. Consider a psych consult if there is evidence of depression, anxiety or irritability. Consider an addiction medicine consult if there is evidence of substance misuse. When to Continue Opioids (a) If the patient has returned to work. (b) If the patient has improved functioning and pain. (Washington, 2002) (Colorado, 2002) (Ontario, 2000) (VA/DoD, 2003) (Maddox-AAPM/APS, 1997) (Wisconsin, 2004) (Warfield, 2004) The long- term use of this medication class is not recommended per the California MTUS unless there documented evidence of benefit with measurable outcome measures and improvement in function. There is no documented significant decrease in objective pain measures such as VAS scores for significant periods of time. There are no objective measures of improvement of function. Therefore all criteria for the ongoing use of opioids have not been met and the request is not medically necessary.