

Case Number:	CM15-0167112		
Date Assigned:	09/11/2015	Date of Injury:	11/21/2013
Decision Date:	10/08/2015	UR Denial Date:	08/03/2015
Priority:	Standard	Application Received:	08/25/2015

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:
 State(s) of Licensure: New Jersey, Alabama, California
 Certification(s)/Specialty: Neurology, Neuromuscular Medicine

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The injured worker is a 47-year-old female who sustained an industrial injury on 11-21-13. A review of the medical records indicates that the injured worker is undergoing treatment for bilateral carpal tunnel syndrome and left shoulder impingement syndrome. Medical records (07-23-15) indicate that the injured worker complains of numbness tingling, cramps and sharp pain in her right hand. The physical exam (07-23-15) reveals painful dorsiflexion of the left wrist. Treatment has included bilateral carpal tunnel release, depomedrol injections to the left shoulder, and 100 physical therapy visits for the left shoulder, as well as medications and psychological counseling. Per the records the left carpal tunnel release was 05-08-15 and has received 8 physical therapy sessions to the left wrist. The original utilization review (08-03-15) non certified the physical therapy to the left wrist as there is no indication as to why the injured worker cannot perform rehabilitation with a home exercise program.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

Physical therapy 2x4 of the left wrist: Upheld

Claims Administrator guideline: Decision based on MTUS Postsurgical Treatment 2009.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Medical Treatment 2009, Section(s): Physical Medicine.

Decision rationale: According to MTUS guidelines, Physical Medicine is "Recommended as indicated below. Passive therapy (those treatment modalities that do not require energy expenditure on the part of the patient) can provide short-term relief during the early phases of pain treatment and are directed at controlling symptoms such as pain, inflammation and swelling and to improve the rate of healing soft tissue injuries. They can be used sparingly with active therapies to help control swelling, pain and inflammation during the rehabilitation process. Active therapy is based on the philosophy that therapeutic exercise and/or activity are beneficial for restoring flexibility, strength, endurance, function, range of motion, and can alleviate discomfort. Active therapy requires an internal effort by the individual to complete a specific exercise or task. This form of therapy may require supervision from a therapist or medical provider such as verbal, visual and/or tactile instruction(s). Patients are instructed and expected to continue active therapies at home as an extension of the treatment process in order to maintain improvement levels. Home exercise can include exercise with or without mechanical assistance or resistance and functional activities with assistive devices. (Colorado, 2002) (Airaksinen, 2006) Patient-specific hand therapy is very important in reducing swelling, decreasing pain, and improving range of motion in CRPS. (Li, 2005) The use of active treatment modalities (e.g., exercise, education, activity modification) instead of passive treatments is associated with substantially better clinical outcomes. In a large case series of patients with low back pain treated by physical therapists, those adhering to guidelines for active rather than passive treatments incurred fewer treatment visits, cost less, and had less pain and less disability. The overall success rates were 64.7% among those adhering to the active treatment recommendations versus 36.5% for passive treatment." (Fritz, 2007). The patient in this case underwent left wrist carpal tunnel release on May 18, 2015. According to the progress report dated June 11, 2015, the patient stated that she felt physical therapy made her left wrist more painful. In addition, post-op physical therapy after carpal tunnel release is recommended for up to 8 sessions for up to 5 weeks after surgery. There is no documentation on the number of physical therapy already been received. Therefore, the request for 8 physical therapy sessions for the left wrist is not medically necessary.