

<b>Case Number:</b>	CM15-0168999		
<b>Date Assigned:</b>	09/04/2015	<b>Date of Injury:</b>	04/28/2010
<b>Decision Date:</b>	10/07/2015	<b>UR Denial Date:</b>	07/23/2015
<b>Priority:</b>	Standard	<b>Application Received:</b>	08/27/2015

### HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:

State(s) of Licensure: North Carolina

Certification(s)/Specialty: Family Practice

### CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The injured worker is a 52 year old male who sustained an industrial lifting injury on 04-28-2010. The injured worker was diagnosed with lumbosacral degenerative intervertebral disc, lumbosacral radiculopathy, lumbosacral radiculitis, chronic pain syndrome, shoulder, and upper arm sprain. The injured worker is status post right shoulder surgery in 2010. Treatment to date has included diagnostic testing with recent lumbar spine magnetic resonance imaging (MRI) in Apr 2015, shoulder surgery, acupuncture therapy, physical therapy, lumbar epidural steroid injection, functional restoration program (FRP), cane and medications. According to the primary treating physician's progress report on July 14, 2015 the injured worker continues to experience low back pain radiating to the left lower extremity into his toes associated with weakness, numbness and tingling of the left lower extremity. Examination demonstrated an antalgic gait favoring the left side with a forward flexed body posture and using a cane for ambulation. Seated straight leg raise was positive on the left at 60 degrees. Range of motion of the hips, knees, ankles and feet were within normal limits. Soft tissue tenderness was noted over the plantar fascia of the left lower extremity with dorsiflexion at 3+ out of 5 on the left and 5 out of 5 on the right. Current medications were listed as Norco 10mg-325mg (taper), Prevacid and Miralax. Treatment plan consists of continuing with walking, exercise and stretching as tolerated, trial of Gabapentin, trial of transcutaneous electrical nerve stimulation (TEN's) unit and the current request for Tramadol 50mg.

### IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

**Tramadol HCL 50mg 1 tablet twice a day #60: Upheld**

**Claims Administrator guideline:** Decision based on MTUS Chronic Pain Treatment Guidelines Opioids Page(s): 91, 77, and 78.

**MAXIMUS guideline:** Decision based on MTUS Chronic Pain Treatment Guidelines opioids Page(s): 76-84.

**Decision rationale:** The California chronic pain medical treatment guidelines section on opioids states for ongoing management: On-Going management, actions should include: (a) Prescriptions from a single practitioner taken as directed and all prescriptions from a single pharmacy. (b) The lowest possible dose should be prescribed to improve pain and function. (c) Office: Ongoing review and documentation of pain relief, functional status, appropriate medication use, and side effects. Pain assessment should include current pain; the least reported pain over the period since last assessment; average pain; intensity of pain after taking the opioid; how long it takes for pain relief; and how long pain relief lasts. Satisfactory response to treatment may be indicated by the patient's decreased pain, increased level of function, or improved quality of life. Information from family members or other caregivers should be considered in determining the patient's response to treatment. The 4 A's for Ongoing Monitoring: Four domains have been proposed as most relevant for ongoing monitoring of chronic pain patients on opioids: pain relief, side effects, physical and psychosocial functioning, and the occurrence of any potentially aberrant (or non-adherent) drug-related behaviors. These domains have been summarized as the "4 A's" (analgesia, activities of daily living, adverse side effects, and aberrant drug taking behaviors). The monitoring of these outcomes over time should affect therapeutic decisions and provide a framework for documentation of the clinical use of these controlled drugs. (Passik, 2000) (d) Home: To aid in pain and functioning assessment, the patient should be requested to keep a pain diary that includes entries such as pain triggers, and incidence of end-of-dose pain. It should be emphasized that using this diary will help in tailoring the opioid dose. This should not be a requirement for pain management. (e) Use of drug screening or inpatient treatment with issues of abuse, addiction, or poor pain control. (f) Documentation of misuse of medications (doctor- shopping, uncontrolled drug escalation, drug diversion). (g) Continuing review of overall situation with regard to non-opioid means of pain control. (h) Consideration of a consultation with a multidisciplinary pain clinic if doses of opioids are required beyond what is usually required for the condition or pain does not improve on opioids in 3 months. Consider a psych consult if there is evidence of depression, anxiety or irritability. Consider an addiction medicine consult if there is evidence of substance misuse. When to Continue Opioids: (a) If the patient has returned to work; (b) If the patient has improved functioning and pain (Washington, 2002) (Colorado, 2002) (Ontario, 2000) (VA/DoD, 2003) (Maddox- AAPM/APS, 1997) (Wisconsin, 2004) (Warfield, 2004). The long-term use of this medication class is not recommended per the California MTUS unless there documented evidence of benefit with measurable outcome measures and improvement in function. There is no documented significant decrease in objective pain measures such as VAS scores for significant periods of time. There are no objective measures of improvement of function. Therefore not all criteria for the ongoing use of opioids have been met and the request is not medically necessary.

