

Case Number:	CM15-0169872		
Date Assigned:	09/10/2015	Date of Injury:	08/08/2001
Decision Date:	10/08/2015	UR Denial Date:	08/24/2015
Priority:	Standard	Application Received:	08/28/2015

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:

State(s) of Licensure: North Carolina

Certification(s)/Specialty: Family Practice

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The injured worker is a 58 year old female, who sustained an industrial injury on 8-8-01. The injured worker has complaints of low back pain with radiation to bilateral buttocks. The documentation noted that the injured worker tried Chantix to stop smoking, it didn't work and she gained 20 pounds. Dietary restrictions and recommendations were reviewed. The documentation noted that a psychology evaluation and treatment for pain related depression, weight gain has been approved and appointment has been scheduled. Palpation of the lumbar facet reveals pain on both the sides at L3-S1 (sacroiliac) region. There is pain noted over the lumbar intervertebral spaces (discs) on palpation. Anterior flexion of lumbar spine is noted to be 40 degrees, anterior lumbar flexion causes pain. Extension of lumbar spine is noted to be 10 degrees and there is pain noted with lumbar extension. The diagnoses have included thoracic or lumbosacral neuritis or radiculitis, unspecified. Treatment to date has included transcutaneous electrical nerve stimulation unit; heating pads; amitriptyline; flexeril; MSContin; norco; Senna; trazodone; facet injections; epidural injections without long lasting relief; physical therapy without relief and acupuncture with improvement. The documentation noted on 7-13-15 the injured worker had reduced soma to zero and is taking flexeril 3 times day for spasms and the flexeril is effective at this frequency. The documentation noted that the injured worker has done well to reduce her medications over the last year we have been treating her that she initially was prescribed 680mg morphine equivalent and has been able to reduce her to 360mg. The original utilization review (8-24-15) partially approved a request for flexeril 10mg #10 (original request for #90) to allow for weaning. The request for MS Contin 100mg #90 was modified to MS Contin 100mg #60 to allow for weaning. The request for Norco 10-325mg #90 was modified to

Norco 10-325mg #60 in order to maintain efficacy during the weaning process.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

Flexeril 10mg #90: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Medical Treatment 2009.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Medical Treatment 2009, Section(s): Muscle relaxants (for pain).

Decision rationale: The California chronic pain medical treatment guidelines section on muscle relaxants states: Recommend non-sedating muscle relaxants with caution as a second-line option for short-term treatment of acute exacerbations in patients with chronic LBP. (Chou, 2007) (Mens, 2005) (Van Tulder, 1998) (Van Tulder, 2003) (Van Tulder, 2006) (Schnitzer, 2004) (See, 2008) Muscle relaxants may be effective in reducing pain and muscle tension, and increasing mobility. However, in most LBP cases, they show no benefit beyond NSAIDs in pain and overall improvement. Also there is no additional benefit shown in combination with NSAIDs. Efficacy appears to diminish over time, and prolonged use of some medications in this class may lead to dependence. (Homik, 2004) (Chou, 2004) This medication is not intended for long-term use per the California MTUS. The medication has not been prescribed for the flare-up of chronic low back pain. This is not an approved use for the medication. For these reasons, criteria for the use of this medication have not been met. Therefore the request is not certified and therefore is not medically necessary.

MS Contin 100mg #90: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Medical Treatment 2009.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Medical Treatment 2009, Section(s): Opioids for chronic pain.

Decision rationale: The California chronic pain medical treatment guidelines section on opioids states for ongoing management: On-Going Management. Actions Should Include: (a) Prescriptions from a single practitioner taken as directed, and all prescriptions from a single pharmacy. (b) The lowest possible dose should be prescribed to improve pain and function. (c) Office: Ongoing review and documentation of pain relief, functional status, appropriate medication use, and side effects. Pain assessment should include: current pain; the least reported pain over the period since last assessment; average pain; intensity of pain after taking the opioid; how long it takes for pain relief; and how long pain relief lasts. Satisfactory response to treatment may be indicated by the patient's decreased pain, increased level of function, or improved quality of life. Information from family members or other caregivers should be

considered in determining the patient's response to treatment. The 4 A's for Ongoing Monitoring: Four domains have been proposed as most relevant for ongoing monitoring of chronic pain patients on opioids: pain relief, side effects, physical and psychosocial functioning, and the occurrence of any potentially aberrant (or non-adherent) drug-related behaviors. These domains have been summarized as the "4 A's" (analgesia, activities of daily living, adverse side effects, and aberrant drug taking behaviors). The monitoring of these outcomes over time should affect therapeutic decisions and provide a framework for documentation of the clinical use of these controlled drugs. (Passik, 2000) (d) Home: To aid in pain and functioning assessment, the patient should be requested to keep a pain diary that includes entries such as pain triggers, and incidence of end-of-dose pain. It should be emphasized that using this diary will help in tailoring the opioid dose. This should not be a requirement for pain management. (e) Use of drug screening or inpatient treatment with issues of abuse, addiction, or poor pain control. (f) Documentation of misuse of medications (doctor-shopping, uncontrolled drug escalation, drug diversion). (g) Continuing review of overall situation with regard to non-opioid means of pain control. (h) Consideration of a consultation with a multidisciplinary pain clinic if doses of opioids are required beyond what is usually required for the condition or pain does not improve on opioids in 3 months. Consider a psych consult if there is evidence of depression, anxiety or irritability. Consider an addiction medicine consult if there is evidence of substance misuse. When to Continue Opioids: (a) If the patient has returned to work; (b) If the patient has improved functioning and pain (Washington, 2002) (Colorado, 2002) (Ontario, 2000) (VA/DoD, 2003) (Maddox-AAPM/APS, 1997) (Wisconsin, 2004) (Warfield, 2004). The long-term use of this medication class is not recommended per the California MTUS unless there documented evidence of benefit with measurable outcome measures and improvement in function. There is no documented significant improvement in VAS scores for significant periods of time. There are no objective measurements of improvement in function. Therefore not all criteria for the ongoing use of opioids have been met and the request is not certified and therefore is not medically necessary.

Norco 10/325mg #90: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Medical Treatment 2009.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Medical Treatment 2009, Section(s): Opioids for chronic pain.

Decision rationale: The California chronic pain medical treatment guidelines section on opioids states for ongoing management: On-Going Management. Actions Should Include: (a) Prescriptions from a single practitioner taken as directed, and all prescriptions from a single pharmacy. (b) The lowest possible dose should be prescribed to improve pain and function. (c) Office: Ongoing review and documentation of pain relief, functional status, appropriate medication use, and side effects. Pain assessment should include: current pain; the least reported pain over the period since last assessment; average pain; intensity of pain after taking the opioid; how long it takes for pain relief; and how long pain relief lasts. Satisfactory response to treatment may be indicated by the patient's decreased pain, increased level of function, or improved quality of life. Information from family members or other caregivers should be

considered in determining the patient's response to treatment. The 4 A's for Ongoing Monitoring: Four domains have been proposed as most relevant for ongoing monitoring of chronic pain patients on opioids: pain relief, side effects, physical and psychosocial functioning, and the occurrence of any potentially aberrant (or non-adherent) drug-related behaviors. These domains have been summarized as the "4 A's" (analgesia, activities of daily living, adverse side effects, and aberrant drug taking behaviors). The monitoring of these outcomes over time should affect therapeutic decisions and provide a framework for documentation of the clinical use of these controlled drugs. (Passik, 2000)

(d) Home: To aid in pain and functioning assessment, the patient should be requested to keep a pain diary that includes entries such as pain triggers, and incidence of end-of-dose pain. It should be emphasized that using this diary will help in tailoring the opioid dose. This should not be a requirement for pain management.

(e) Use of drug screening or inpatient treatment with issues of abuse, addiction, or poor pain control.

(f) Documentation of misuse of medications (doctor-shopping, uncontrolled drug escalation, drug diversion).

(g) Continuing review of overall situation with regard to non-opioid means of pain control.

(h) Consideration of a consultation with a multidisciplinary pain clinic if doses of opioids are required beyond what is usually required for the condition or pain does not improve on opioids in 3 months. Consider a psych consult if there is evidence of depression, anxiety or irritability. Consider an addiction medicine consult if there is evidence of substance misuse.

When to Continue Opioids: (a) If the patient has returned to work; (b) If the patient has improved functioning and pain (Washington, 2002) (Colorado, 2002) (Ontario, 2000) (VA/DoD, 2003) (Maddox-AAPM/APS, 1997) (Wisconsin, 2004) (Warfield, 2004). The long-term use of this medication class is not recommended per the California MTUS unless there documented evidence of benefit with measurable outcome measures and improvement in function. There is no documented significant improvement in VAS scores for significant periods of time. There are no objective measurements of improvement in function. Therefore not all criteria for the ongoing use of opioids have been met and the request is not certified and therefore is not medically necessary.