

<b>Case Number:</b>	CM15-0180851		
<b>Date Assigned:</b>	09/22/2015	<b>Date of Injury:</b>	11/09/1998
<b>Decision Date:</b>	11/02/2015	<b>UR Denial Date:</b>	09/11/2015
<b>Priority:</b>	Standard	<b>Application Received:</b>	09/14/2015

### HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:

State(s) of Licensure: North Carolina

Certification(s)/Specialty: Family Practice

### CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The injured worker is a 58-year-old male who sustained an industrial injury on November 9, 1998. Diagnoses have included persistent right side greater than left-sided neck pain and chronic low back pain. Documented treatment includes cervical spinal fusion at C4-C6 in 2000, Medial branch blocks C4-C6 March 6, 2015, and medications including Percocet, Cymbalta, Lidoderm patches, Prevacid, and Ambien stated in the February 12, 2015 progress note to enable him to perform activities of daily living and exercise. The injured worker continues to report neck, back and upper extremity pain with some paresthesias down his right arm. Physician noted positive cervical facet maneuvers and tenderness over the paraspinal regions. The treating physician's plan of care includes Lidoderm patches with 2 refills, but this was denied on September 1, 2015. Current work status is stated as "sedentary work only" but it is not clarified if he is working at present.

### IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

**Lidoderm patch 5%, #30 with 2 refills:** Upheld

**Claims Administrator guideline:** Decision based on MTUS Chronic Pain Medical Treatment 2009.

**MAXIMUS guideline:** Decision based on MTUS Chronic Pain Medical Treatment 2009, Section(s): Topical Analgesics.

**Decision rationale:** The California chronic pain medical treatment guidelines section on topical lidocaine states: Lidocaine Indication: Neuropathic pain Recommended for localized peripheral pain after there has been evidence of a trial of first-line therapy (tri-cyclic or SNRI anti-depressants or an AED such as gabapentin or Lyrica). The FDA for neuropathic pain has designated topical lidocaine, in the formulation of a dermal patch (Lidoderm) for orphan status. Lidoderm is also used off-label for diabetic neuropathy. No other commercially approved topical formulations of lidocaine (whether creams, lotions or gels) are indicated for neuropathic pain. Non-dermal patch formulations are generally indicated as local anesthetics and anti-pruritics. Further research is needed to recommend this treatment for chronic neuropathic pain disorders other than post-herpetic neuralgia. Formulations that do not involve a dermal-patch system are generally indicated as local anesthetics and anti-pruritics. In February 2007 the FDA notified consumers and healthcare professionals of the potential hazards of the use of topical lidocaine. Those at particular risk were individuals that applied large amounts of this substance over large areas, left the products on for long periods of time, or used the agent with occlusive dressings. Systemic exposure was highly variable among patients. Only FDA-approved products are currently recommended. (Argoff, 2006) (Dworkin, 2007) (Khaliq-Cochrane, 2007) (Knotkova, 2007) (Lexi-Comp, 2008) Non-neuropathic pain: Not recommended. There is only one trial that tested 4% lidocaine for treatment of chronic muscle pain. The results showed there was no superiority over placebo. (Scudds, 1995) This medication is recommended for localized peripheral pain. The patient does have back and neck pain, however the patient has no documented failure of all first line agents indicated for the treatment of neuropathic pain as outlined above. Therefore, criteria as set forth by the California MTUS as outlined above have not been met and the request is not medically necessary.