

<b>Case Number:</b>	CM15-0182759		
<b>Date Assigned:</b>	09/23/2015	<b>Date of Injury:</b>	05/09/2013
<b>Decision Date:</b>	11/03/2015	<b>UR Denial Date:</b>	08/20/2015
<b>Priority:</b>	Standard	<b>Application Received:</b>	09/17/2015

### HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:  
 State(s) of Licensure: California, District of Columbia, Maryland  
 Certification(s)/Specialty: Anesthesiology, Pain Management

### CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The injured worker is a 55 year old female who sustained an industrial injury on 5-9-13. The injured worker reported left knee discomfort. A review of the medical records indicates that the injured worker is undergoing treatments for lumbosacral sprain strain, lumbar muscle spasm, lumbar disc protrusion, left knee sprain strain, left knee medial meniscus tear. Medical records dated 6-18-15 indicate pain rated at 4 out of 10 on average. Provider documentation dated 5-12-15 noted the work status as return to modified work 5-13-15. Treatment has included Naproxen since at least June of 2015, Ibuprofen since at least June of 2015, and Menthoderm topical analgesic cream since at least June of 2015, left knee radiographic studies, left knee magnetic resonance imaging and acupuncture treatment. Physical examination dated 6-18-15 was notable for left posterior tibial tendon and left Achilles tendon with pain upon range of motion and "antalgic limp favoring the right side." The original utilization review (8-20-15) denied a request for Prilosec-Omeprazole 20 milligrams quantity of 90 prescribed 07-27-2015, Nabutone 500 milligrams quantity of 60 prescribed 07-27-2015, and Cyclobenzaprine 10 milligrams quantity of 60 prescribed 07-27-2015.

### IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

**Prilosec/Omeprazole 20mg #90 prescribed 07/27/2015: Upheld**

**Claims Administrator guideline:** Decision based on MTUS Chronic Pain Medical Treatment 2009.

**MAXIMUS guideline:** Decision based on MTUS Chronic Pain Medical Treatment 2009, Section(s): NSAIDs, GI symptoms & cardiovascular risk.

**Decision rationale:** In the treatment of dyspepsia secondary to NSAID therapy, the MTUS recommends stopping the NSAID, switching to a different NSAID, or considering the use of an H2-receptor antagonist or a PPI. The MTUS Chronic Pain Medical Treatment Guidelines recommend the use of proton pump inhibitors in conjunction with NSAIDs in situations in which the patient is at risk for gastrointestinal events including: (1) age > 65 years; (2) history of peptic ulcer, GI bleeding or perforation; (3) concurrent use of ASA, corticosteroids, and/or an anticoagulant; or (4) high dose/multiple NSAID (e.g., NSAID + low-dose ASA). CPMTG guidelines further specify: "Recommendations: Patients with no risk factor and no cardiovascular disease: Non-selective NSAIDs OK (e.g, ibuprofen, naproxen, etc.) Patients at intermediate risk for gastrointestinal events and no cardiovascular disease:(1) A non-selective NSAID with either a PPI (Proton Pump Inhibitor, for example, 20 mg omeprazole daily) or misoprostol (200 ug four times daily) or (2) a Cox-2 selective agent. Long-term PPI use (> 1 year) has been shown to increase the risk of hip fracture (adjusted odds ratio 1.44). Patients at high risk for gastrointestinal events with no cardiovascular disease: A Cox-2 selective agent plus a PPI if absolutely necessary. Patients at high risk of gastrointestinal events with cardiovascular disease: If GI risk is high the suggestion is for a low-dose Cox-2 plus low dose Aspirin (for cardioprotection) and a PPI. If cardiovascular risk is greater than GI risk the suggestion is naproxyn plus low-dose aspirin plus a PPI. (Laine, 2006) (Scholmerich, 2006) (Nielsen, 2006) (Chan, 2004) (Gold, 2007) (Laine, 2007)" As there is no documentation of peptic ulcer, GI bleeding or perforation, or cardiovascular disease in the records available for my review, the injured worker's risk for gastrointestinal events is low, as such, medical necessity cannot be affirmed.

**Nabutone 500mg #60 prescribed 07/27/2015: Upheld**

**Claims Administrator guideline:** Decision based on MTUS Chronic Pain Medical Treatment 2009.

**MAXIMUS guideline:** Decision based on MTUS Chronic Pain Medical Treatment 2009, Section(s): NSAIDs (non-steroidal anti-inflammatory drugs).

**Decision rationale:** With regard to the use of NSAIDs for chronic low back pain, the MTUS CPMTG states "Recommended as an option for short-term symptomatic relief. A Cochrane review of the literature on drug relief for low back pain (LBP) suggested that NSAIDs were no more effective than other drugs such as acetaminophen, narcotic analgesics, and muscle relaxants. The review also found that NSAIDs had more adverse effects than placebo and acetaminophen but fewer effects than muscle relaxants and narcotic analgesics. In addition, evidence from the review suggested that no one NSAID, including COX-2 inhibitors, was clearly more effective than another." The documentation submitted for review indicates that the injured

worker has been using NSAID medications ibuprofen and naproxen since at least 6/2015. As NSAIDs are only recommended for short-term symptomatic relief, the request is not medically necessary.

**Cyclobenzaprine 10mg #60 prescribed 07/27/2015: Upheld**

**Claims Administrator guideline:** Decision based on MTUS Chronic Pain Medical Treatment 2009.

**MAXIMUS guideline:** Decision based on MTUS Chronic Pain Medical Treatment 2009, Section(s): Muscle relaxants (for pain).

**Decision rationale:** With regard to muscle relaxants, the MTUS CPMTG states: "Recommend non-sedating muscle relaxants with caution as a second-line option for short-term treatment of acute exacerbations in patients with chronic LBP. (Chou, 2007) (Mens, 2005) (Van Tulder, 1998) (van Tulder, 2003) (Van Tulder, 2006) (Schnitzer, 2004) (See, 2008) Muscle relaxants may be effective in reducing pain and muscle tension, and increasing mobility. However, in most LBP cases, they show no benefit beyond NSAIDs in pain and overall improvement." Regarding Cyclobenzaprine: "Recommended for a short course of therapy. Limited, mixed-evidence does not allow for a recommendation for chronic use. Cyclobenzaprine is a skeletal muscle relaxant and a central nervous system depressant with similar effects to tricyclic antidepressants (e.g. amitriptyline). Cyclobenzaprine is more effective than placebo in the management of back pain, although the effect is modest and comes at the price of adverse effects." Per p41 of the MTUS guidelines the effect is greatest in the first 4 days of treatment, suggesting that shorter courses may be better. Treatment is recommended for the treatment of acute spasm limited to a maximum of 2-3 weeks. Per the medical records, the patient is not being treated for an acute exacerbation of chronic back pain, but for left knee pain. The requested treatment is not medically necessary.