

Case Number:	CM15-0184595		
Date Assigned:	10/15/2015	Date of Injury:	08/05/2004
Decision Date:	11/23/2015	UR Denial Date:	08/31/2015
Priority:	Standard	Application Received:	09/19/2015

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:
 State(s) of Licensure: North Carolina
 Certification(s)/Specialty: Family Practice

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

This is a 44 year old female who sustained an industrial injury on 8-5-2004. A review of the medical records indicates that the injured worker is undergoing treatment for cervical disc disease, lumbar strain, lumbar radiculitis, bilateral shoulder sprain, post surgery headaches, bilateral sacroiliitis, anxiety and stress, cervicogenic headaches and migraine headaches. Medical records (9-17-2015) indicate ongoing pain from the neck going down into both shoulders, elbows and hands. The injured worker also complained of frequent migraines. She complained of low back pain with radiation down the left leg. The physical exam (9-17-2015) revealed tenderness to palpation of the cervical spine. Spurling's test caused radicular symptoms to the upper extremities. There was stiffness and tightness on deep palpation at L4-L5 as well as bilateral, posterior, superior iliac spine. Treatment has included cervical spine surgery and medications. The injured worker has been prescribed Motrin since at least 6-1-2015. Previous progress reports list Fenoprofen. Other medications (7-6-2015) included Tylenol #3, Prilosec and Lunesta. The original Utilization Review (UR) (8-31-2015) denied requests for Motrin and Botox.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

Motrin 800mg #60: Overturned

Claims Administrator guideline: Decision based on MTUS Chronic Pain Medical Treatment 2009.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Medical Treatment 2009, Section(s): NSAIDs (non-steroidal anti-inflammatory drugs).

Decision rationale: The California chronic pain medical treatment guidelines section on NSAID therapy states: Recommended at the lowest dose for the shortest period in patients with moderate to severe pain. Acetaminophen may be considered for initial therapy for patients with mild to moderate pain, and in particular, for those with gastrointestinal, cardiovascular or renovascular risk factors. NSAIDs appear to be superior to acetaminophen, particularly for patients with moderate to severe pain. There is no evidence to recommend one drug in this class over another based on efficacy. In particular, there appears to be no difference between traditional NSAIDs and COX-2 NSAIDs in terms of pain relief. The main concern of selection is based on adverse effects. COX-2 NSAIDs have fewer GI side effects at the risk of increased cardiovascular side effects, although the FDA has concluded that long-term clinical trials are best interpreted to suggest that cardiovascular risk occurs with all NSAIDs and is a class effect (with naproxyn being the safest drug). There is no evidence of long-term effectiveness for pain or function. (Chen, 2008) (Laine, 2008) Back Pain - Chronic low back pain: Recommended as an option for short-term symptomatic relief. A Cochrane review of the literature on drug relief for low back pain (LBP) suggested that NSAIDs were no more effective than other drugs such as acetaminophen, narcotic analgesics, and muscle relaxants. The review also found that NSAIDs had more adverse effects than placebo and acetaminophen but fewer effects than muscle relaxants and narcotic analgesics. In addition, evidence from the review suggested that no one NSAID, including COX-2 inhibitors, was clearly more effective than another. (Roelofs-Cochrane, 2008) See also Anti-inflammatory medications. Neuropathic pain: There is inconsistent evidence for the use of these medications to treat long term neuropathic pain, but they may be useful to treat breakthrough and mixed pain conditions such as osteoarthritis (and other nociceptive pain) in with neuropathic pain. This medication is recommended for the shortest period of time and at the lowest dose possible. The dosing of this medication is within the California MTUS guideline recommendations. The definition of shortest period possible is not clearly defined in the California MTUS. Therefore the request is medically necessary.

One (1) Botox injection: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Medical Treatment 2009.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Medical Treatment 2009, Section(s): Botulinum toxin (Botox Myobloc).

Decision rationale: The California chronic pain medical treatment guidelines section on botulism toxin states: Not generally recommended for chronic pain disorders, but recommended for cervical dystonia. Not recommended for the following: tension-type headache; migraine headache; fibromyositis; chronic neck pain; myofascial pain syndrome; & trigger point

injections. Several recent studies have found no statistical support for the use of Botulinum toxin A (BTXA) for any of the following: The evidence is mixed for migraine headaches. This RCT found that both botulinum toxin type A (BoNTA) and divalproex sodium (DVPX) significantly reduced disability associated with migraine, and BoNTA had a favorable tolerability profile compared with DVPX. (Blumenfeld, 2008) In this RCT of episodic migraine patients, low-dose injections of BoNTA into the frontal, temporal, and/or glabellar muscle regions were not more effective than placebo. (Saper, 2007) Botulinum neurotoxin is probably ineffective in episodic migraine and chronic tension-type headache (Level B). (Naumann, 2008)- Myofascial analgesic pain relief as compared to saline. (Qerama, 2006)- Use as a specific treatment for myofascial cervical pain as compared to saline. (Ojala, 2006) (Ferrante, 2005) (Wheeler, 1998)- Injection in myofascial trigger points as compared to dry needling or local anesthetic injections. (Kamanli, 2005) (Graboski, 2005). Recent systematic reviews have stated that current evidence does not support the use of BTX-A trigger point injections for myofascial pain. (Ho, 2006) Or for mechanical neck disease (as compared to saline). (Peloso-Cochrane, 2006) A recent study that has found statistical improvement with the use of BTX-A compared to saline. Study patients had at least 10 trigger points and no patient in the study was allowed to take an opioid in the 4 weeks prior to treatment. (Gobel, 2006) Recommended: cervical dystonia, a condition that is not generally related to workers-compensation injuries (also known as spasmodic torticollis), and is characterized as a movement disorder of the nuchal muscles, characterized by tremor or by tonic posturing of the head in a rotated, twisted, or abnormally flexed or extended position or some combination of these positions. When treated with BTX-B, high antigenicity limits long-term efficacy. Botulinum toxin A injections provide more objective and subjective benefit than trihexyphenidyl or other anticholinergic drugs to patients with cervical dystonia. Recommended: chronic low back pain, if a favorable initial response predicts subsequent responsiveness, as an option in conjunction with a functional restoration program. Some additional new data suggests that it may be effective for low back pain. (Jabbari, 2006) (Ney, 2006) Botulinum neurotoxin may be considered for low back pain (Level C). (Naumann, 2008) The requested medication is usually only indicated in the treatment of cervical dystonia. Per the California MTUS it does not have the indication in the treatment of other diagnosis. The patient does not have this diagnosis but rather neck pain and cervicogenic headaches. Therefore the request is not medically necessary.