

<b>Case Number:</b>	CM15-0188644		
<b>Date Assigned:</b>	09/30/2015	<b>Date of Injury:</b>	09/26/2014
<b>Decision Date:</b>	11/12/2015	<b>UR Denial Date:</b>	09/15/2015
<b>Priority:</b>	Standard	<b>Application Received:</b>	09/24/2015

### HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:  
 State(s) of Licensure: California, District of Columbia, Maryland  
 Certification(s)/Specialty: Anesthesiology, Pain Management

### CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The injured worker is a 52 year old female who sustained an industrial injury on 9-26-2014. A review of medical records indicates the injured worker is being treated for pain in joint of lower leg. Medical records dated 8-31-2015 noted right knee pain. She rates the pain is a 6 out of 10. She is able to walk for 4 blocks and stand for 3 hours. Physical examination noted range of motion to the right knee was restricted. There was tenderness to palpation over the medial joint line and patella. Treatment has included gabapentin, ibuprofen, meloxicam, and baclofen. A prescription was written for Cyclobenzaprine, Lidopro, and Protonix. Utilization review form dated 9-15-2015 noncertified Cyclobenzaprine 7.5mg #60, Protonix 20mg #60, and Lidopro 4% ointment time one tube.

### IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

**Cyclobenzaprine 7.5mg quantity 60:** Upheld

**Claims Administrator guideline:** Decision based on MTUS Chronic Pain Medical Treatment 2009.

**MAXIMUS guideline:** Decision based on MTUS Chronic Pain Medical Treatment 2009, Section(s): Muscle relaxants (for pain).

**Decision rationale:** With regard to muscle relaxants, the MTUS CPMTG states: "Recommend non-sedating muscle relaxants with caution as a second-line option for short-term treatment of acute exacerbations in patients with chronic LBP. (Chou, 2007) (Mens, 2005) (Van Tulder, 1998) (van Tulder, 2003) (van Tulder, 2006) (Schnitzer, 2004) (See, 2008) Muscle relaxants may be effective in reducing pain and muscle tension, and increasing mobility. However, in most LBP cases, they show no benefit beyond NSAIDs in pain and overall improvement." Regarding Cyclobenzaprine: "Recommended for a short course of therapy. Limited, mixed-evidence does not allow for a recommendation for chronic use. Cyclobenzaprine is a skeletal muscle relaxant and a central nervous system depressant with similar effects to tricyclic antidepressants (e.g. amitriptyline). Cyclobenzaprine is more effective than placebo in the management of back pain, although the effect is modest and comes at the price of adverse effects." Per p41 of the MTUS guidelines, the effect is greatest in the first 4 days of treatment, suggesting that shorter courses may be better. Treatment is recommended for the treatment of acute spasm limited to a maximum of 2-3 weeks. The patient is not being treated for an acute exacerbation of chronic back pain, so the requested treatment is not medically necessary.

**Pantoprazole 20mg quantity 60:** Upheld

**Claims Administrator guideline:** The Claims Administrator did not base their decision on the MTUS. Decision based on Non-MTUS Citation Official Disability Guidelines, Proton Pump Inhibitors.

**MAXIMUS guideline:** Decision based on MTUS Chronic Pain Medical Treatment 2009, Section(s): NSAIDs, GI symptoms & cardiovascular risk. Decision based on Non-MTUS Citation Official Disability Guidelines (ODG) Pain, Proton Pump Inhibitors.

**Decision rationale:** In the treatment of dyspepsia secondary to NSAID therapy, the MTUS recommends stopping the NSAID, switching to a different NSAID, or considering the use of an H2-receptor antagonist or a PPI. The MTUS Chronic Pain Medical Treatment Guidelines recommend the use of proton pump inhibitors in conjunction with NSAIDs in situations in which the patient is at risk for gastrointestinal events including: (1) age > 65 years; (2) history of peptic ulcer, GI bleeding or perforation; (3) concurrent use of ASA, corticosteroids, and/or an anticoagulant; or (4) high dose/multiple NSAID (e.g., NSAID + low-dose ASA). CPMTG guidelines further specify: "Recommendations: Patients with no risk factor and no cardiovascular disease: Non-selective NSAIDs OK (e.g, ibuprofen, naproxen, etc.). Patients at intermediate risk for gastrointestinal events and no cardiovascular disease: (1) A non-selective NSAID with either a PPI (Proton Pump Inhibitor, for example, 20 mg omeprazole daily) or misoprostol (200 g four times daily); or (2) a Cox-2 selective agent. Long-term PPI use (> 1 year) has been shown to increase the risk of hip fracture (adjusted odds ratio 1.44). Patients at high risk for gastrointestinal events with no cardiovascular disease: A Cox-2 selective agent plus a PPI if absolutely necessary. Patients at high risk of gastrointestinal events with cardiovascular disease: If GI risk is high the suggestion is for a low-dose Cox-2 plus low dose Aspirin (for cardioprotection) and a PPI. If cardiovascular risk is greater than GI risk, the suggestion is

naproxyn plus low-dose aspirin plus a PPI. (Laine, 2006) (Scholmerich, 2006) (Nielsen, 2006) (Chan, 2004) (Gold, 2007) (Laine, 2007)" Per ODG TWC, "many prescribers believe that this class of drugs is innocuous, but much information is available to demonstrate otherwise. A trial of omeprazole or lansoprazole is recommended before Nexium therapy. The other PPIs, Protonix, Dexilant, and Aciphex, should also be second-line." As there is no documentation of peptic ulcer, GI bleeding or perforation, or cardiovascular disease in the records available for my review, the injured worker's risk for gastrointestinal events is low, as such, medical necessity cannot be affirmed. Furthermore, as noted per the guidelines, Protonix is a second-line medication. The medical records do not establish whether the patient has failed attempts at first line PPIs, such as omeprazole or lansoprazole, which should be considered prior to prescribing a second line PPI such as Protonix. The request is not medically necessary.

**Lidopro 4% Ointment one tube: Upheld**

**Claims Administrator guideline:** Decision based on MTUS Chronic Pain Medical Treatment 2009.

**MAXIMUS guideline:** Decision based on MTUS Chronic Pain Medical Treatment 2009, Section(s): Topical Analgesics.

**Decision rationale:** Regarding the use of multiple medications, MTUS p60 states "Only one medication should be given at a time, and interventions that are active and passive should remain unchanged at the time of the medication change. A trial should be given for each individual medication. Analgesic medications should show effects within 1 to 3 days, and the analgesic effect of antidepressants should occur within 1 week. A record of pain and function with the medication should be recorded. (Mens, 2005) The recent AHRQ review of comparative effectiveness and safety of analgesics for osteoarthritis concluded that each of the analgesics was associated with a unique set of benefits and risks, and no currently available analgesic was identified as offering a clear overall advantage compared with the others." Therefore, it would be optimal to trial each medication individually. LidoPro contains capsaicin, lidocaine, menthol, methyl salicylate. Per MTUS p 112 with regard to capsaicin, "Indications: There are positive randomized studies with capsaicin cream in patients with osteoarthritis, fibromyalgia, and chronic non-specific back pain, but it should be considered experimental in very high doses. Although topical capsaicin has moderate to poor efficacy, it may be particularly useful (alone or in conjunction with other modalities) in patients whose pain has not been controlled successfully with conventional therapy." Methyl salicylate may have an indication for chronic pain in this context. Per MTUS p105, "Recommended. Topical salicylate (e.g., Ben-Gay, methyl salicylate) is significantly better than placebo in chronic pain. (Mason-BMJ, 2004)" However, the other ingredients in LidoPro are not indicated. The preponderance of evidence indicates that overall, this medication is not medically necessary. Regarding topical lidocaine, MTUS states (p112) "Neuropathic pain Recommended for localized peripheral pain after there has been evidence of a trial of first-line therapy (tri-cyclic or SNRI anti-depressants or an AED such as gabapentin or Lyrica). Non-neuropathic pain: Not recommended. There is only one trial that tested 4% lidocaine for treatment of chronic muscle pain. The results showed there was no superiority over placebo. (Scudds, 1995)" The documentation submitted for review does not contain evidence of trial of first-line therapy to support the use of topical lidocaine. LidoPro topical lotion contains

menthol. The CA MTUS, ODG, National Guidelines Clearinghouse, and ACOEM provide no evidence-based recommendations regarding the topical application of menthol. It is the opinion of this IMR reviewer that a lack of endorsement, a lack of mention, inherently implies a lack of recommendation, or a status equivalent to "not recommended". Since menthol is not medically indicated, then the overall product is not indicated per MTUS as outlined below. Note the statement on page 111: Any compounded product that contains at least one drug (or drug class) that is not recommended is not recommended. The request is not medically necessary.