

Case Number:	CM15-0189393		
Date Assigned:	10/01/2015	Date of Injury:	11/20/2009
Decision Date:	12/03/2015	UR Denial Date:	09/08/2015
Priority:	Standard	Application Received:	09/25/2015

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:

State(s) of Licensure: California

Certification(s)/Specialty: Emergency Medicine

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The injured worker is a 54 year old male, who sustained an industrial injury on November 20, 2009. He reported injury to his low back and bilateral shoulders. The injured worker was diagnosed as having low back pain, lumbar radiculopathy and status post bilateral shoulder arthroscopies. Treatment to date has included physical therapy, diagnostic studies and medication. On August 25, 2015, the injured worker complained of constant lower back pain with radiation through the back, sides of his legs, knees and feet accompanied by numbness and tingling. There was a constant, aching to sharp pain across his lower back with intermittent pain radiating down the left leg to the foot. He noted numbness and tingling in his left leg. He complained of constant mid low back pain rated as a 5-6 on a 1-10 pain scale intermittently localized in the lumbosacral area. Physical examination of the lumbosacral area showed tenderness and spasm in the paralumbar area from L1-S1 bilaterally. Mild tenderness was noted in the greater sciatic notch area on the right side and mild tenderness over the sacrococcygeal area. Straight leg raise elicited pain at about 60 degrees on the left and the right. The treatment plan included lumbar epidural steroid series, cyclobenzaprine, hydrocodone, Naprosyn, omeprazole and a follow-up visit. On September 8, 2015, utilization review denied a request for Cyclobenzaprine 7.5mg #90, Hydrocodone 10-325mg #120, Naprosyn 375mg #30 and Omeprazole 20mg #30.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

Cyclobenzaprine 7.5mg #90: Upheld

Claims Administrator guideline: The Claims Administrator did not cite any medical evidence for its decision.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Medical Treatment 2009, Section(s): Muscle relaxants (for pain).

Decision rationale: The request is for the use of a muscle relaxant to aid in pain relief. The MTUS guidelines state that the use of a medication in this class is indicated as a second-line option for short-term treatment of acute exacerbations of low back pain. Muscle relaxants may be effective in reducing pain and muscle tension, which can increase mobility. However, in most LBP cases, they show no benefit beyond NSAIDs in pain improvement. Efficacy appears to diminish over time, and prolonged use may lead to dependence. (Homik, 2004) Due to inadequate documentation of a recent acute exacerbation and poor effectiveness for chronic long-term use, the request is not medically necessary.

Hydrocodone 10/325mg #120: Upheld

Claims Administrator guideline: The Claims Administrator did not cite any medical evidence for its decision.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Medical Treatment 2009, Section(s): Opioids, criteria for use.

Decision rationale: The request is for the use of a medication in the opioid class. The MTUS guidelines state that for ongoing treatment with a pharmaceutical in this class, certain requirements are necessary. This includes not only adequate pain control, but also functional improvement. Four domains have been proposed for management of patients on opioids. This includes pain relief, side effects, physical and psychosocial functioning, and the occurrence of any potentially aberrant drug-related behaviors. As part of the pain treatment agreement, it is advised that "Refills are limited, and will only occur at appointments". In this case, there is inadequate documentation of persistent functional improvement seen. As such, the request is not medically necessary. All opioid medications should be titrated down slowly in order to prevent a significant withdrawal syndrome.

Naprosyn 375mg #30: Upheld

Claims Administrator guideline: The Claims Administrator did not cite any medical evidence for its decision.

MAXIMUS guideline: The Expert Reviewer did not base their decision on the MTUS. Decision based on Non-MTUS Citation Official Disability Guidelines (ODG) Pain (Chronic)/NSAIDs (non-steroidal anti-inflammatory drugs).

Decision rationale: The request is for the use of a medication in the NSAID class. The ODG state the following regarding this topic: Specific recommendations: Osteoarthritis (including knee and hip): Recommended at the lowest dose for the shortest period in patients with moderate to severe pain. Acetaminophen may be considered for initial therapy for patients with mild to moderate pain, and in particular, for those with gastrointestinal, cardiovascular or renovascular risk factors. NSAIDs appear to be superior to acetaminophen, particularly for patients with moderate to severe pain. There is no evidence to recommend one drug in this class over another based on efficacy. In particular, there appears to be no difference between traditional NSAIDs and COX-2 NSAIDs in terms of pain relief. The main concern of selection is based on adverse effects. COX-2 NSAIDs have fewer GI side effects at the risk of increased cardiovascular side effects, although the FDA has concluded that long-term clinical trials are best interpreted to suggest that cardiovascular risk occurs with all NSAIDs and is a class effect (with naproxen being the safest drug). There is no evidence of long-term effectiveness for pain or function. (Chen, 2008) (Laine, 2008) Back Pain - Acute low back pain & acute exacerbations of chronic pain: Recommended as a second-line treatment after acetaminophen. In general, there is conflicting to negative evidence that NSAIDs are more effective than acetaminophen for acute LBP. (Van Tulder, 2006) (Hancock, 2007) For patients with acute low back pain with sciatica a recent Cochrane review (including three heterogeneous randomized controlled trials) found no differences in treatment with NSAIDs vs. placebo. In patients with axial low back pain this same review found that NSAIDs were not more effective than acetaminophen for acute low-back pain, and that acetaminophen had fewer side effects. (Roelofs-Cochrane, 2008) The addition of NSAIDs or spinal manipulative therapy does not appear to increase recovery in patients with acute low back pain over that received with acetaminophen treatment and advice from their physician. (Hancock, 2007) Back Pain - Chronic low back pain: Recommended as an option for short-term symptomatic relief. A Cochrane review of the literature on drug relief for low back pain (LBP) suggested that NSAIDs were no more effective than other drugs such as acetaminophen, narcotic analgesics, and muscle relaxants. The review also found that NSAIDs had more adverse effects than placebo and acetaminophen but fewer effects than muscle relaxants and narcotic analgesics. In addition, evidence from the review suggested that no one NSAID, including COX-2 inhibitors, was clearly more effective than another. (Roelofs-Cochrane, 2008) See also Anti-inflammatory medications. Neuropathic pain: There is inconsistent evidence for the use of these medications to treat long-term neuropathic pain, but they may be useful to treat breakthrough pain and mixed pain conditions such as osteoarthritis (and other nociceptive pain) in patients with neuropathic pain. (Namaka, 2004) (Gore, 2006) See NSAIDs, GI symptoms & cardiovascular risk; NSAIDs, hypertension and renal function; & Medications for acute pain (analgesics). Besides the above well-documented side effects of NSAIDs, there are other less well-known effects of NSAIDs, and the use of NSAIDs has been shown to possibly delay and hamper healing in all the soft tissues, including muscles, ligaments, tendons, and cartilage. (Maroon, 2006) The risks of NSAIDs in older patients, which include increased cardiovascular risk and gastrointestinal toxicity, may outweigh the benefits of these medications. (AGS, 2009) As stated above, acetaminophen would be considered first-line treatment for chronic pain. In this case, the continued use of an NSAID is not supported. This is secondary to inadequate documentation of functional improvement benefit seen. Also, the duration of use places the patient at risk for gastrointestinal and cardiovascular side-effects. In

addition, it is known that use of NSAIDs delays the healing of soft tissue including ligaments, tendons, and cartilage. As such, the request is not medically necessary.

Omeprazole 20mg #30: Upheld

Claims Administrator guideline: The Claims Administrator did not cite any medical evidence for its decision.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Medical Treatment 2009, Section(s): NSAIDs, GI symptoms & cardiovascular risk.

Decision rationale: The request is for the use of a medication in the class of a proton pump inhibitor. It is indicated for patients with peptic ulcer disease. It can also be used as a preventative measure in patients taking non-steroidal anti-inflammatories for chronic pain. Unfortunately, they do have certain side effects including gastrointestinal disease. The MTUS guidelines states that patients who are classified as intermediate or high risk, should be treated prophylactically. Criteria for risk are as follows: "(1) age > 65 years; (2) history of peptic ulcer, GI bleeding or perforation; (3) concurrent use of ASA, corticosteroids, and/or an anticoagulant; or (4) high dose/multiple NSAID (e.g., NSAID + low-dose ASA)." Due to the fact the patient does not meet to above stated criteria, the request for use is not medically necessary.