

Case Number:	CM15-0195166		
Date Assigned:	10/08/2015	Date of Injury:	05/25/2012
Decision Date:	11/23/2015	UR Denial Date:	10/05/2015
Priority:	Standard	Application Received:	10/05/2015

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:
 State(s) of Licensure: California, District of Columbia, Maryland
 Certification(s)/Specialty: Anesthesiology, Pain Management

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The injured worker is a 46 year old female who sustained an industrial injury 05-25-12. A review of the medical records reveals the injured worker is undergoing treatment for cervical and lumbar spine sprain-strain, persistent right knee, right wrist, and right shoulder complaints, as well as left hand numbness and tingling. Medical records (09-22-15) reveal the injured worker complains of shoulder and upper extremity pain. The pain is not rated. The physical exam (09-22-15) reveals diminished range of motion in the right shoulder as well as tenderness on the volar aspect. Lumbar and cervical spine range of motion is also noted to be diminished. Tenderness is noted in the paraspinal muscles. Prior treatment includes right knee, right carpal tunnel syndrome, and right shoulder surgeries; home exercise program, injections, and medications. The original utilization review (10-05-15) non certified the request for Flexeril 10mg #30 and Norco 10/325 #60. The documentation supports that he injured worker has been on Norco since at least 06-16-15 and Flexeril since at least 08-31-15.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

Flexeril 10mg Qty: 30.00: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Medical Treatment 2009, Section(s): Muscle relaxants (for pain).

MAXIMUS guideline: Decision based on MTUS Chronic Pain Medical Treatment 2009, Section(s): Muscle relaxants (for pain).

Decision rationale: With regard to muscle relaxants, the MTUS CPMTG states: "Recommend non-sedating muscle relaxants with caution as a second-line option for short-term treatment of acute exacerbations in patients with chronic LBP. (Chou, 2007) (Mens, 2005) (Van Tulder, 1998) (van Tulder, 2003) (van Tulder, 2006) (Schnitzer, 2004) (See, 2008) Muscle relaxants may be effective in reducing pain and muscle tension, and increasing mobility. However, in most LBP cases, they show no benefit beyond NSAIDs in pain and overall improvement." Regarding Cyclobenzaprine: "Recommended for a short course of therapy. Limited, mixed-evidence does not allow for a recommendation for chronic use. Cyclobenzaprine is a skeletal muscle relaxant and a central nervous system depressant with similar effects to tricyclic antidepressants (e.g. amitriptyline). Cyclobenzaprine is more effective than placebo in the management of back pain, although the effect is modest and comes at the price of adverse effects." Per p41 of the MTUS guidelines the effect is greatest in the first 4 days of treatment, suggesting that shorter courses may be better. Treatment is recommended for the treatment of acute spasm limited to a maximum of 2-3 weeks. UDS that evaluate for cyclobenzaprine can provide additional data on whether the injured worker is compliant, however in this case there is no UDS testing for cyclobenzaprine. The documentation submitted for review indicates that the injured worker has been using this medication since at least 10/2014. There is no documentation of the patient's specific functional level or percent improvement with treatment with cyclobenzaprine. As it is recommended only for short-term use, medical necessity cannot be affirmed, therefore not medically necessary.

Norco 10/325mg Qty: 60.00: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Medical Treatment 2009, Section(s): Opioids, pain treatment agreement.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Medical Treatment 2009, Section(s): Opioids, criteria for use.

Decision rationale: Per MTUS Chronic Pain Medical Treatment Guidelines p78 regarding on-going management of opioids Four domains have been proposed as most relevant for ongoing monitoring of chronic pain patients on opioids: Pain relief, side effects, physical and psychosocial functioning, and the occurrence of any potentially aberrant (or nonadherent) drug related behaviors. These domains have been summarized as the 4 A's (Analgesia, activities of daily living, adverse side effects, and any aberrant drug-taking behaviors). The monitoring of these outcomes over time should affect therapeutic decisions and provide a framework for documentation of the clinical use of these controlled drugs. Review of the available medical records reveals no documentation to support the medical necessity of Norco or any documentation addressing the '4 A's' domains, which is a recommended practice for the on-going management of opioids. Specifically, the notes do not appropriately review and document pain

relief, functional status improvement, appropriate medication use, or side effects. The MTUS considers this list of criteria for initiation and continuation of opioids in the context of efficacy required to substantiate medical necessity, and they do not appear to have been addressed by the treating physician in the documentation available for review. Furthermore, efforts to rule out aberrant behavior (e.g. CURES report, UDS, opiate agreement) are necessary to assure safe usage and establish medical necessity. There is no documentation comprehensively addressing this concern in the records available for my review. As MTUS recommends discontinuing opioids if there is no overall improvement in function, medical necessity cannot be affirmed, therefore is not medically necessary.