

<b>Case Number:</b>	CM15-0196322		
<b>Date Assigned:</b>	10/09/2015	<b>Date of Injury:</b>	11/15/2012
<b>Decision Date:</b>	11/18/2015	<b>UR Denial Date:</b>	09/30/2015
<b>Priority:</b>	Standard	<b>Application Received:</b>	10/06/2015

### HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:

State(s) of Licensure: North Carolina

Certification(s)/Specialty: Family Practice

### CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The injured worker is a 63 year old male who sustained an industrial injury on 11-15-2012. Medical records indicated the worker was treated for shoulder pain, left shoulder chronic tendinitis-bursitis status post decompression, cervicgia and lumbago. The worker rates her pain as 5 on a scale of 10 with medications, 8 on a scale of 0-10 without medications, and states there are no new problems or side effect with the medications. Quality of sleep is poor. Current medications include Ibuprofen and omeprazole. On examination, his cervical spine reveals straightening of the spine with loss of normal cervical lordosis. Range of motion is restricted by pain. There is tenderness, hypertonicity and spasm noted on the cervical paravertebral muscles bilaterally. On examination of the thoracic spine, there is hyper tonicity, spasm, and a tight muscle band bilaterally. The left shoulder has no limitation in range of motion. Hawkins test is positive, Neer test is positive. Empty cans test is positive. There are no gastrointestinal complaints. A request for authorization was submitted for Omeprazole 20 mg #90 with 2 refills. A utilization review decision 09-30-2015 gave modified approval of Omeprazole with none of the requested refills.

### IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

**Omeprazole 20 mg #90 with 2 refills:** Upheld

**Claims Administrator guideline:** Decision based on MTUS Chronic Pain Medical Treatment 2009, Section(s): NSAIDs (non-steroidal anti-inflammatory drugs).

**MAXIMUS guideline:** Decision based on MTUS Chronic Pain Medical Treatment 2009, Section(s): NSAIDs, GI symptoms & cardiovascular risk.

**Decision rationale:** The California chronic pain medical treatment guidelines section on NSAID therapy and proton pump inhibitors (PPI) states: Recommend with precautions as indicated below. Clinicians should weight the indications for NSAIDs against both GI and cardiovascular risk factors. Determine if the patient is at risk for gastrointestinal events: (1) age > 65 years; (2) history of peptic ulcer, GI bleeding or perforation; (3) concurrent use of ASA, corticosteroids, and/or an anticoagulant; or (4) high dose/multiple NSAID (e.g., NSAID + low-dose ASA). Recent studies tend to show that H. Pylori does not act synergistically with NSAIDS to develop gastro duodenal lesions. Recommendations; Patients with no risk factor and no cardiovascular disease: Non-selective NSAIDs OK. (e.g., ibuprofen, naproxen, etc.). Patients at intermediate risk for gastrointestinal events and no cardiovascular disease: (1) A non-selective NSAID with either a PPI (Proton Pump Inhibitor, for example, 20 mg omeprazole daily) or misoprostol (200 g four times daily) or (2) a Cox-2 selective agent. Long-term PPI use (> 1 year) has been shown to increase the risk of hip fracture (adjusted odds ratio 1.44). Patients at high risk for gastrointestinal events with no cardiovascular disease: A Cox-2 selective agent plus a PPI if absolutely necessary. There is no documentation provided that places this patient at intermediate or high risk that would justify the use of a PPI. There is no mention of current gastrointestinal or cardiovascular disease. For these reasons the criteria set forth above per the California MTUS for the use of this medication has not been met. Therefore, the request is not medically necessary.