

Case Number:	CM15-0199969		
Date Assigned:	10/15/2015	Date of Injury:	04/18/2001
Decision Date:	11/24/2015	UR Denial Date:	10/09/2015
Priority:	Standard	Application Received:	10/12/2015

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:

State(s) of Licensure: Montana

Certification(s)/Specialty: Preventive Medicine, Occupational Medicine

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The injured worker is a 66 year old female who suffered an industrial injury on 4-18-2001. The diagnoses included cervical spine fusion 2008, post-operative cervical spine syndrome with spondylosis without myelopathy, cervical disc degeneration, thoracic disc degeneration and thoracic pain. On 9-24-2015 the treating provider reported cervical and low back pain rated on average 8 out of 10. She reported high levels of pain located in the cervical spine across the shoulders and between the shoulder blades. On examination the cervical range of motion was reduced by 50% and elevated cervical and thoracic muscle spasticity without triggering. The provider noted no neurologic complaints or significant neurologic examination findings. He requested re-imaging of the cervical spine with prior MRI in early 2013. He also noted she may have an occult thoracic injury or disc herniation however, no radiographs are noted. Prior treatment included prior cervical fusion C3-5 in 2008, failed left cervical radiofrequency ablation in 8-2015 and 9-2015, medication, physical therapy, home exercise program, acupuncture, chiropractic and trigger point injections. Request for Authorization date was 10-2-2015. The Utilization Review on 10-9-2015 determined non-certification for MRI of cervical spine with and without contrast and MRI of the thoracic spine with and without contrast.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

MRI of cervical spine with and without contrast: Upheld

Claims Administrator guideline: The Claims Administrator did not base their decision on the MTUS. Decision based on Non-MTUS Citation Official Disability Guidelines, MRI.

MAXIMUS guideline: Decision based on MTUS Neck and Upper Back Complaints 2004, Section(s): Special Studies. Decision based on Non-MTUS Citation Official Disability Guidelines (ODG) Neck, Magnetic resonance imaging (MRI).

Decision rationale: The MTUS states that for most patients presenting with true neck or upper back problems, special studies are not needed unless a three- or four-week period of conservative care and observation fails to improve symptoms. Most patients improve quickly, provided any red-flag conditions are ruled out. Criteria for ordering imaging studies are: emergence of a red flag, physiologic evidence of tissue insult or neurologic dysfunction, failure to progress in a strengthening program intended to avoid surgery and clarification of the anatomy prior to an invasive procedure. Physiologic evidence may be in the form of definitive neurologic findings on physical examination, electrodiagnostic studies, laboratory tests, or bone scans. Unequivocal findings that identify specific nerve compromise on the neurologic examination are sufficient evidence to warrant imaging studies if symptoms persist. When the neurologic examination is less clear, however, further physiologic evidence of nerve dysfunction can be obtained before ordering an imaging study. Electromyography (EMG), and nerve conduction velocities (NCV), including H-reflex tests, may help identify subtle focal neurologic dysfunction in patients with neck or arm symptoms, or both, lasting more than three or four weeks. The assessment may include sensory-evoked potentials (SEPs) if spinal stenosis or spinal cord myelopathy is suspected. If physiologic evidence indicates tissue insult or nerve impairment, consider a discussion with a consultant regarding next steps, including the selection of an imaging test to define a potential cause (magnetic resonance imaging [MRI] for neural or other soft tissue, compute tomography [CT] for bony structures). Additional studies may be considered to further define problem areas. The recent evidence indicates cervical disk annular tears may be missed on MRIs. The clinical significance of such a finding is unclear, as it may not correlate temporally or anatomically with symptoms. Reliance on imaging studies alone to evaluate the source of neck or upper back symptoms carries a significant risk of diagnostic confusion (false-positive test results) because it's possible to identify a finding that was present before symptoms began and, therefore, has no temporal association with the symptoms. The ODG guidelines state that magnetic resonance imaging (MRI) is not recommended except for indications list below. Patients who are alert, have never lost consciousness, are not under the influence of alcohol and/or drugs, have no distracting injuries, have no cervical tenderness, and have no neurologic findings, do not need imaging. Patients who do not fall into this category should have a three-view cervical radiographic series followed by computed tomography (CT). In determining whether or not the patient has ligamentous instability, magnetic resonance imaging (MRI) is the procedure of choice, but MRI should be reserved for patients who have clear-cut neurologic findings and those suspected of ligamentous instability. Repeat MRI is not routinely recommended, and should be reserved for a significant change in symptoms and/or findings suggestive of significant pathology (eg, tumor, infection, fracture, neurocompression, recurrent disc herniation). (Anderson, 2000) (ACR, 2002) See also ACR Appropriateness Criteria. MRI imaging studies are valuable when physiologic evidence indicates tissue insult or nerve

impairment or potentially serious conditions are suspected like tumor, infection, and fracture, or for clarification of anatomy prior to surgery. MRI is the test of choice for patients who have had prior back surgery. (Bigos, 1999) (Bey, 1998) (Volle, 2001) (Singh, 2001) (Colorado, 2001) For the evaluation of the patient with chronic neck pain, plain radiographs (3-view: anteroposterior, lateral, open mouth) should be the initial study performed. Patients with normal radiographs and neurologic signs or symptoms should undergo magnetic resonance imaging. If there is a contraindication to the magnetic resonance examination such as a cardiac pacemaker or severe claustrophobia, computed tomography myelography, preferably using spiral technology and multiplanar reconstruction is recommended. (Daffner, 2000) (Bono, 2007) Indications for imaging MRI (magnetic resonance imaging): Chronic neck pain (= after 3 months conservative treatment), radiographs normal, neurologic signs or symptoms present; Neck pain with radiculopathy if severe or progressive neurologic deficit; Chronic neck pain, radiographs show spondylosis, neurologic signs or symptoms present; Chronic neck pain, radiographs show old trauma, neurologic signs or symptoms present; Chronic neck pain, radiographs show bone or disc margin destruction; Suspected cervical spine trauma, neck pain, clinical findings suggest ligamentous injury (sprain), radiographs and/or CT "normal"; Known cervical spine trauma: equivocal or positive plain films with neurological deficit; Upper back/thoracic spine trauma with neurological deficit. In this case there is no documentation of neurologic deficit or significant progression of symptoms. The treating physician has not indicated concern about any "red flag" conditions. The request for MRI of cervical spine with and without contrast is not consistent with the MTUS and ODG guidelines and is not medically necessary.

MRI of the thoracic spine with and without contrast: Upheld

Claims Administrator guideline: The Claims Administrator did not base their decision on the MTUS. Decision based on Non-MTUS Citation Official Disability Guidelines, MRI.

MAXIMUS guideline: Decision based on MTUS Neck and Upper Back Complaints 2004, Section(s): Special Studies. Decision based on Non-MTUS Citation Official Disability Guidelines (ODG) Neck, Magnetic resonance imaging (MRI).

Decision rationale: The MTUS states that for most patients presenting with true neck or upper back problems, special studies are not needed unless a three- or four-week period of conservative care and observation fails to improve symptoms. Most patients improve quickly, provided any red-flag conditions are ruled out. Criteria for ordering imaging studies are: emergence of a red flag, physiologic evidence of tissue insult or neurologic dysfunction, failure to progress in a strengthening program intended to avoid surgery and clarification of the anatomy prior to an invasive procedure. Physiologic evidence may be in the form of definitive neurologic findings on physical examination, electrodiagnostic studies, laboratory tests, or bone scans. Unequivocal findings that identify specific nerve compromise on the neurologic examination are sufficient evidence to warrant imaging studies if symptoms persist. When the neurologic examination is less clear, however, further physiologic evidence of nerve dysfunction can be obtained before ordering an imaging study. Electromyography (EMG), and nerve conduction velocities (NCV), including H-reflex tests, may help identify subtle focal neurologic dysfunction in patients with neck or arm symptoms, or both, lasting more than three or four weeks. The assessment may include sensory-evoked potentials (SEPs) if spinal stenosis or spinal cord myelopathy is

suspected. If physiologic evidence indicates tissue insult or nerve impairment, consider a discussion with a consultant regarding next steps, including the selection of an imaging test to define a potential cause (magnetic resonance imaging [MRI] for neural or other soft tissue, compute tomography [CT] for bony structures). Additional studies may be considered to further define problem areas. The recent evidence indicates cervical disk annular tears may be missed on MRIs. The clinical significance of such a finding is unclear, as it may not correlate temporally or anatomically with symptoms. Reliance on imaging studies alone to evaluate the source of neck or upper back symptoms carries a significant risk of diagnostic confusion (false-positive test results) because it's possible to identify a finding that was present before symptoms began and, therefore, has no temporal association with the symptoms. The ODG guidelines state that magnetic resonance imaging (MRI) is not recommended except for indications list below. Patients who are alert, have never lost consciousness, are not under the influence of alcohol and/or drugs, have no distracting injuries, have no cervical tenderness, and have no neurologic findings, do not need imaging. Patients who do not fall into this category should have a three-view cervical radiographic series followed by computed tomography (CT). In determining whether or not the patient has ligamentous instability, magnetic resonance imaging (MRI) is the procedure of choice, but MRI should be reserved for patients who have clear-cut neurologic findings and those suspected of ligamentous instability. Repeat MRI is not routinely recommended, and should be reserved for a significant change in symptoms and/or findings suggestive of significant pathology (eg, tumor, infection, fracture, neurocompression, recurrent disc herniation). (Anderson, 2000) (ACR, 2002) See also ACR Appropriateness Criteria. MRI imaging studies are valuable when physiologic evidence indicates tissue insult or nerve impairment or potentially serious conditions are suspected like tumor, infection, and fracture, or for clarification of anatomy prior to surgery. MRI is the test of choice for patients who have had prior back surgery. (Bigos, 1999) (Bey, 1998) (Volle, 2001) (Singh, 2001) (Colorado, 2001) For the evaluation of the patient with chronic neck pain, plain radiographs (3-view: anteroposterior, lateral, open mouth) should be the initial study performed. Patients with normal radiographs and neurologic signs or symptoms should undergo magnetic resonance imaging. If there is a contraindication to the magnetic resonance examination such as a cardiac pacemaker or severe claustrophobia, computed tomography myelography, preferably using spiral technology and multiplanar reconstruction is recommended. (Daffner, 2000) (Bono, 2007) Indications for imaging MRI (magnetic resonance imaging): Chronic neck pain (= after 3 months conservative treatment), radiographs normal, neurologic signs or symptoms present; Neck pain with radiculopathy if severe or progressive neurologic deficit; Chronic neck pain, radiographs show spondylosis, neurologic signs or symptoms present; Chronic neck pain, radiographs show old trauma, neurologic signs or symptoms present; Chronic neck pain, radiographs show bone or disc margin destruction; Suspected cervical spine trauma, neck pain, clinical findings suggest ligamentous injury (sprain), radiographs and/or CT "normal"; Known cervical spine trauma: equivocal or positive plain films with neurological deficit; Upper back/thoracic spine trauma with neurological deficit. In this case there is no documentation of neurologic deficit or significant progression of symptoms. The treating physician has not indicated concern about any "red flag" conditions. The request for MRI of the thoracic spine with and without contrast is not consistent with the MTUS and ODG guidelines and is not medically necessary.