

Case Number:	CM15-0200039		
Date Assigned:	10/15/2015	Date of Injury:	01/12/2007
Decision Date:	11/24/2015	UR Denial Date:	09/28/2015
Priority:	Standard	Application Received:	10/12/2015

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:
 State(s) of Licensure: California, Oregon, Washington
 Certification(s)/Specialty: Orthopedic Surgery

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The injured worker is a 54 year old male who sustained an industrial injury on 01-12-2007. A review of the medical records indicated that the injured worker is undergoing treatment for cervicgia, cervical sprain and strain and cervicogenic headaches. The injured worker is status post lumbar fusion (no date documented). According to the treating physician's progress report on 09-10-2015, the injured worker continues to experience headaches without associated symptoms and right sided neck pain. The injured worker also reported low back pain which is the current worst pain. Average pain was noted as 8 out of 10 on the pain scale along with poor sleep quality. Examination of the cervical spine was brief noting right sided neck pain and crepitus. Prior treatments specifically for the cervical spine were not noted. Current medications were listed as Hydrocodone, Hysingla ER, Percocet and Ambien. Urine drug screening tests over the past 5 months were reported as consistent within the 09-10-2015 report. Treatment plan consists of continuing medication regimen, regular home exercise on an ongoing and regular basis, Naprosyn and esomeprazole prescriptions, consultation for shoulder and sacroiliac area and the current request for right cervical C2-3, C3-4 and C4-5 medial branch blocks. On 09-28-2015, the Utilization Review determined the request for right cervical C2-3, C3-4, C4-5 medial branch blocks was not medically necessary.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

Right cervical C2-3, C3-4, C4-5 medial branch blocks: Upheld

Claims Administrator guideline: Decision based on MTUS Neck and Upper Back Complaints 2004, Section(s): Initial Care. Decision based on Non-MTUS Citation Official Disability Guidelines (ODG), Neck and Upper Back, Facet joint diagnostic blocks.

MAXIMUS guideline: Decision based on MTUS Neck and Upper Back Complaints 2004, Section(s): Initial Care, Follow-up Visits. Decision based on Non-MTUS Citation Official Disability Guidelines (ODG) neck section / facet joint diagnostic blocks.

Decision rationale: CA MTUS/ACOEM Chapter 8, Neck and Upper Back Complaints, Table 8-8, page 181, does not recommend facet injection of corticosteroids or diagnostic blocks in the cervical spine. As the guidelines do not recommend facet blocks, the determination is for non-certification. ODG-TWC, neck section notes that facet joint diagnostic blocks are recommended prior to facet neurotomy (a procedure that is considered “under study”). Diagnostic blocks are performed with the anticipation that if successful, treatment may proceed to facet neurotomy at the diagnosed levels. Criteria for the use of diagnostic blocks for facet nerve pain: Clinical presentation should be consistent with facet joint pain, signs & symptoms. 1. One set of diagnostic medial branch blocks is required with a response of 70%. The pain response should be approximately 2 hours for Lidocaine. 2. Limited to patients with cervical pain that is non-radicular and at no more than two levels bilaterally. 3. There is documentation of failure of conservative treatment (including home exercise, PT and NSAIDs) prior to the procedure for at least 4-6 weeks. 4. No more than 2 joint levels are injected in one session (see above for medial branch block levels). 5. Recommended volume of no more than 0.5 cc of injectate is given to each joint, with recent literature suggesting a volume of 0.25 cc to improve diagnostic accuracy. 6. No pain medication from home should be taken for at least 4 hours prior to the diagnostic block and for 4 to 6 hours afterward. 7. Opioids should not be given as a "sedative" during the procedure. 8. The use of IV sedation may be grounds to negate the results of a diagnostic block, and should only be given in cases of extreme anxiety. 9. The patient should document pain relief with an instrument such as a VAS scale, emphasizing the importance of recording the maximum pain relief and maximum duration of pain. The patient should also keep medication use and activity logs to support subjective reports of better pain control. 10. Diagnostic facet blocks should not be performed in patients in whom a surgical procedure is anticipated. 11. Diagnostic facet blocks should not be performed in patients who have had a previous fusion procedure at the planned injection level. 12. It is currently not recommended to perform facet blocks on the same day of treatment as epidural steroid injections or stellate ganglion blocks or sympathetic blocks or trigger point injections as this may lead to improper diagnosis or unnecessary treatment. As the guidelines do not recommend facet blocks, the determination is for non-certification. There is no documentation if a failure of appropriate nonoperative treatment. No more than 2 levels should be done at one time.