

Case Number:	CM15-0208124		
Date Assigned:	10/27/2015	Date of Injury:	11/17/2007
Decision Date:	12/14/2015	UR Denial Date:	09/28/2015
Priority:	Standard	Application Received:	10/22/2015

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:

State(s) of Licensure: Massachusetts

Certification(s)/Specialty: Anesthesiology, Pain Management

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The injured worker is a 47 year old female who sustained an industrial injury on 9-28-15. She is not working. The medical records indicate that the injured worker has been treated for back pain; sciatica; gastritis due to medication; mild L4-5 radiculitis; obesity; right ankle strain, resolved. She currently (9-15-15) complains of ankle and low back pain. On physical exam there was tenderness at L3-4, L4-5, spasms on the right lumbar area, trigger points L3, L4, L5 and right sciatic, sacroiliac joint tenderness, decreased range of motion by 75%, reduced sensation in the foot and calf, deep tendon reflexes reduced; there was tenderness of the medial and lateral right ankle. The 4-7-15 note indicated a pain level of 7 out of 10. Treatments to date include medication: Flexeril, Lyrica with benefit, hydrocodone, Lidoderm, omeprazole; trigger point injections with benefit for acute flare ups; cervical epidural steroid injection (1-8-15); lumbar epidural steroid injection with benefit (50%). Prior physical therapy documentation was not found. The request for authorization dated 9-17-15 was for physical therapy to the lumbar spine 3 times a week for 4 weeks. On 9-28-15 Utilization Review non-certified the request for physical therapy to the lumbar spine 3 times a week for 4 weeks.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

Physical therapy for the lumbar spine 3 times a week for 4 weeks, quantity: 12 sessions:
Upheld

Claims Administrator guideline: Decision based on MTUS Low Back Complaints 2004, Section(s): Summary.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Medical Treatment 2009, Section(s): Physical Medicine.

Decision rationale: Physical Medicine is recommended as indicated below. Passive therapy (those treatment modalities that do not require energy expenditure on the part of the injured worker) can provide short term relief during the early phases of pain treatment and are directed at controlling symptoms such as pain, inflammation and swelling and to improve the rate of healing soft tissue injuries. They can be used sparingly with active therapies to help control swelling, pain and inflammation during the rehabilitation process. Active therapy is based on the philosophy that therapeutic exercise and/or activity are beneficial for restoring flexibility, strength, endurance, function, range of motion, and can alleviate discomfort. Active therapy requires an internal effort by the individual to complete a specific exercise or task. This form of therapy may require supervision from a therapist or medical provider such as verbal, visual and/or tactile instruction(s). Injured workers are instructed and expected to continue active therapies at home as an extension of the treatment process in order to maintain improvement levels. Home exercise can include exercise with or without mechanical assistance or resistance and functional activities with assistive devices. (Colorado, 2002) (Airaksinen, 2006) Injured worker-specific hand therapy is very important in reducing swelling, decreasing pain, and improving range of motion in CRPS. (Li, 2005) The use of active treatment modalities (e.g., exercise, education, activity modification) instead of passive treatments is associated with substantially better clinical outcomes. In a large case series of injured workers with low back pain treated by physical therapists, those adhering to guidelines for active rather than passive treatments incurred fewer treatment visits, cost less, and had less pain and less disability. The overall success rates were 64.7% among those adhering to the active treatment recommendations versus 36.5% for passive treatment. (Fritz, 2007) Physical Medicine Guidelines Allow for fading of treatment frequency (from up to 3 visits per week to 1 or less), plus active self-directed home Physical Medicine. Myalgia and myositis, unspecified (ICD9 729.1): 9-10 visits over 8 weeks. Neuralgia, neuritis, and radiculitis, unspecified (ICD9 729.2) 8-10 visits over 4 weeks. Reflex sympathetic dystrophy (CRPS) (ICD9 337.2): 24 visits over 16 weeks. According to the documents available for review, the requested number of PT sessions is in excess to the guidelines as set forth in the MTUS. Therefore, at this time, the requirements for treatment have not been met and not medically necessary.