

Case Number:	CM15-0024080		
Date Assigned:	02/13/2015	Date of Injury:	06/28/2006
Decision Date:	04/02/2015	UR Denial Date:	02/06/2015
Priority:	Standard	Application Received:	02/09/2015

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:

State(s) of Licensure: Hawaii, California, Iowa

Certification(s)/Specialty: Preventive Medicine, Occupational Medicine

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

This 63 year old female sustained an industrial injury on 6/28/06. She subsequently reports chronic low back pain with radiation to the lower extremities. Diagnoses include lumbago. Treatment to date has included acupuncture, injections, physical therapy, chiropractic care and prescription medications. On 2/6/15, Utilization Review non-certified a request for Norco 10/325mg QTY: 170.00 with 2 Refills and Soma 350mg QTY: 40.00 with 2 Refills. The Norco 10/325mg QTY: 170.00 with 2 Refills and Soma 350mg QTY: 40.00 with 2 Refills were denied based on MTUS Chronic Pain guidelines.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

Norco 10/325mg QTY: 170.00: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Page(s): 78-80, 91, 124.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Opioids Page(s): 74-96. Decision based on Non-MTUS Citation Official Disability Guidelines (ODG)

Neck and Upper Back (Acute and Chronic), Low Back - Lumbar & Thoracic (Acute & Chronic), Opioids, Pain.

Decision rationale: ODG does not recommend the use of opioids for neck and low back pain "except for short use for severe cases, not to exceed 2 weeks." The patient has exceeded the 2 week recommended treatment length for opioid usage. MTUS does not discourage use of opioids past 2 weeks, but does state that "ongoing review and documentation of pain relief, functional status, appropriate medication use, and side effects. Pain assessment should include: current pain; the least reported pain over the period since last assessment; average pain; intensity of pain after taking the opioid; how long it takes for pain relief; and how long pain relief lasts. Satisfactory response to treatment may be indicated by the patient's decreased pain, increased level of function, or improved quality of life." The treating physician does not fully document the least reported pain over the period since last assessment, intensity of pain after taking opioid, pain relief, increased level of function, or improved quality of life. As such, the request for Norco 10/325mg QTY: 170.00 is not medically necessary.

Refill of norco 10/325mg QTY: 170.00: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Page(s): 78-80, 91, 124.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Opioids Page(s): 74-96. Decision based on Non-MTUS Citation Official Disability Guidelines (ODG) Neck and Upper Back (Acute and Chronic), Low Back - Lumbar & Thoracic (Acute & Chronic), Opioids, Pain.

Decision rationale: ODG does not recommend the use of opioids for neck and low back pain "except for short use for severe cases, not to exceed 2 weeks." The patient has exceeded the 2 week recommended treatment length for opioid usage. MTUS does not discourage use of opioids past 2 weeks, but does state that "ongoing review and documentation of pain relief, functional status, appropriate medication use, and side effects. Pain assessment should include: current pain; the least reported pain over the period since last assessment; average pain; intensity of pain after taking the opioid; how long it takes for pain relief; and how long pain relief lasts. Satisfactory response to treatment may be indicated by the patient's decreased pain, increased level of function, or improved quality of life." The treating physician does not fully document the least reported pain over the period since last assessment, intensity of pain after taking opioid, pain relief, increased level of function, or improved quality of life. Refills of Norco #170 would allow for at least 2 months of this medication without any interim evaluation by the health care provider. Refills are not appropriate in this case. As such, the request for refill of Norco 10/325mg QTY: 170.00 is not medically necessary.

Refill of norco 10/325mg QTY: 170.00: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Page(s): 78-80, 91, 124.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Opioids Page(s): 74-96. Decision based on Non-MTUS Citation Official Disability Guidelines (ODG) Neck and Upper Back (Acute and Chronic), Low Back - Lumbar & Thoracic (Acute & Chronic), Opioids, Pain.

Decision rationale: ODG does not recommend the use of opioids for neck and low back pain "except for short use for severe cases, not to exceed 2 weeks." The patient has exceeded the 2 week recommended treatment length for opioid usage. MTUS does not discourage use of opioids past 2 weeks, but does state that "ongoing review and documentation of pain relief, functional status, appropriate medication use, and side effects. Pain assessment should include: current pain; the least reported pain over the period since last assessment; average pain; intensity of pain after taking the opioid; how long it takes for pain relief; and how long pain relief lasts. Satisfactory response to treatment may be indicated by the patient's decreased pain, increased level of function, or improved quality of life." The treating physician does not fully document the least reported pain over the period since last assessment, intensity of pain after taking opioid, pain relief, increased level of function, or improved quality of life. Refills of Norco #170 would allow for at least 2 months of this medication without any interim evaluation by the health care provider. Refills are not appropriate in this case. As such, the request for refill of Norco 10/325mg QTY: 170.00 is not medically necessary.

Soma 350mg QTY: 40.00: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Page(s): 65.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Carisoprodol (Soma) and Muscle relaxants (for pain) Page(s): 29, 63-66. Decision based on Non-MTUS Citation Official Disability Guidelines (ODG) Chronic Pain, Soma (Carisoprodol).

Decision rationale: MTUS states regarding Carisoprodol, "Not recommended. This medication is not indicated for long-term use. Carisoprodol is a commonly prescribed, centrally acting skeletal muscle relaxant whose primary active metabolite is meprobamate (a schedule-IV controlled substance). Carisoprodol is now scheduled in several states but not on a federal level. It has been suggested that the main effect is due to generalized sedation and treatment of anxiety. Abuse has been noted for sedative and relaxant effects. In regular abusers the main concern is the accumulation of meprobamate. Carisoprodol abuse has also been noted in order to augment or alter effects of other drugs." ODG States that Soma is "Not recommended. This medication is FDA-approved for symptomatic relief of discomfort associated with acute pain in musculoskeletal conditions as an adjunct to rest and physical therapy (AHFS, 2008). This medication is not indicated for long-term use." Per the treatment notes, Soma is to be taken 1 pill at night. The request would allow for at least 40 days of this medication. The medical records do not indicate the extenuating circumstances necessary to deviate from 'long term' usage of this medication, which is high risk for abuse. As such, the request for Soma 350mg QTY: 40.00 is not medically necessary.

Refill of soma 350mg QTY: 40.00: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Page(s): 65.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Carisoprodol (Soma) and Muscle relaxants (for pain) Page(s): 29, 63-66. Decision based on Non-MTUS Citation Official Disability Guidelines (ODG) Chronic Pain, Soma (Carisoprodol).

Decision rationale: MTUS states regarding Carisoprodol, "Not recommended. This medication is not indicated for long-term use. Carisoprodol is a commonly prescribed, centrally acting skeletal muscle relaxant whose primary active metabolite is meprobamate (a schedule-IV controlled substance). Carisoprodol is now scheduled in several states but not on a federal level. It has been suggested that the main effect is due to generalized sedation and treatment of anxiety. Abuse has been noted for sedative and relaxant effects. In regular abusers the main concern is the accumulation of meprobamate. Carisoprodol abuse has also been noted in order to augment or alter effects of other drugs."ODG States that Soma is "Not recommended. This medication is FDA-approved for symptomatic relief of discomfort associated with acute pain in musculoskeletal conditions as an adjunct to rest and physical therapy (AHFS, 2008). This medication is not indicated for long-term use." Per the treatment notes, Soma is to be taken 1 pill at night. This refill request would allow for at least 40 additional days of this medication. The medical records do not indicate the extenuating circumstances necessary to deviate from 'long term' usage of this medication, which is high risk for abuse. As such, the request for Refill of Soma 350mg QTY: 40.00 is not medically necessary.

Refill of soma 350mg QTY: 40.00: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Page(s): 65.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Carisoprodol (Soma) and Muscle relaxants (for pain) Page(s): 29, 63-66. Decision based on Non-MTUS Citation Official Disability Guidelines (ODG) Chronic Pain, Soma (Carisoprodol).

Decision rationale: MTUS states regarding Carisoprodol, "Not recommended. This medication is not indicated for long-term use. Carisoprodol is a commonly prescribed, centrally acting skeletal muscle relaxant whose primary active metabolite is meprobamate (a schedule-IV controlled substance). Carisoprodol is now scheduled in several states but not on a federal level. It has been suggested that the main effect is due to generalized sedation and treatment of anxiety. Abuse has been noted for sedative and relaxant effects. In regular abusers the main concern is the accumulation of meprobamate. Carisoprodol abuse has also been noted in order to augment or alter effects of other drugs."ODG States that Soma is "Not recommended. This medication is FDA-approved for symptomatic relief of discomfort associated with acute pain in musculoskeletal conditions as an adjunct to rest and physical therapy (AHFS, 2008). This medication is not indicated for long-term use."Per the treatment notes, Soma is to be taken 1 pill at night. This refill request would allow for at least 40 additional days of this medication. The medical records do not indicate the extenuating circumstances necessary to deviate from 'long

term' usage of this medication, which is high risk for abuse. As such, the request for Refill of Soma 350mg QTY: 40.00 is not medically necessary.