

Case Number:	CM15-0024720		
Date Assigned:	02/17/2015	Date of Injury:	03/31/2003
Decision Date:	04/02/2015	UR Denial Date:	01/28/2015
Priority:	Standard	Application Received:	02/09/2015

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:

State(s) of Licensure: Massachusetts

Certification(s)/Specialty: Anesthesiology, Pain Management

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The injured worker is a 51 year old female who sustained an industrial injury on 03/31/03. She reports chronic low back pain. Treatments to date include medications. Diagnoses include low back pain, chronic pain syndrome, and lumbago. In a progress note dated 01/16/15 the treating provider recommends continued treatment with Norco and Carisoprodol. On 01/28/15 Utilization Review non-certified the Norco and Carisoprodol, citing MTUS guidelines.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

Pharmacy purchase of Carisoprodol 350 mg: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Muscle Relaxants.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Carisoprodol (Soma) Page(s): 29.

Decision rationale: According to the MTUS, Carisoprodol is not recommended. This medication is not indicated for long-term use. Carisoprodol is a commonly prescribed, centrally

acting skeletal muscle relaxant whose primary active metabolite is meprobamate (a schedule-IV controlled substance). Carisoprodol is now scheduled in several states but not on a federal level. It has been suggested that the main effect is due to generalized sedation and treatment of anxiety. Abuse has been noted for sedative and relaxant effects. In regular abusers the main concern is the accumulation of meprobamate. Carisoprodol abuse has also been noted in order to augment or alter effects of other drugs. This includes the following: (1) increasing sedation of benzodiazepines or alcohol; (2) use to prevent side effects of cocaine; (3) use with tramadol to produce relaxation and euphoria; (4) as a combination with hydrocodone, an effect that some abusers claim is similar to heroin (referred to as a Las Vegas Cocktail); & (5) as a combination with codeine (referred to as Soma Coma). (Reeves, 1999)(Reeves, 2001) (Reeves, 2008) (Schears, 2004) There was a 300% increase in numbers of emergency room episodes related to carisoprodol from 1994 to 2005. (DHSS, 2005) Intoxication appears to include subdued consciousness, decreased cognitive function, and abnormalities of the eyes, vestibular function, appearance, gait and motor function. Intoxication includes the effects of both carisoprodol and meprobamate, both of which act on different neurotransmitters. (Bramness, 2007) (Bramness, 2004) A withdrawal syndrome has been documented that consists of insomnia, vomiting, tremors, muscle twitching, anxiety, and ataxia when abrupt discontinuation of large doses occurs. This is similar to withdrawal from meprobamate. (Reeves, 2007) (Reeves, 2004) There is little research in terms of weaning of high dose carisoprodol and there is no standard treatment regimen for injured workers with known dependence. Most treatment includes treatment for symptomatic complaints of withdrawal. Another option is to switch to phenobarbital to prevent withdrawal with subsequent tapering. A maximum dose of phenobarbital is 500 mg/day and the taper is 30 mg/day with a slower taper in an out-injured worker setting. Tapering should be individualized for each injured worker. (Boothby, 2003) For more information and references, see Muscle relaxants. See also Weaning of medications. Therefore, at this time, the requirements for treatment have not been met and medical necessity has not been established.

Norco #75 with two refills: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Opioids for Chronic Pain.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Opioids, On-Going Management Page(s): 74-97.

Decision rationale: According to the MTUS Chronic Pain Medical Treatment Guidelines section on Opioids, On-Going Management, p 74-97, (a) Prescriptions from a single practitioner taken as directed, and all prescriptions from a single pharmacy. (b) The lowest possible dose should be prescribed to improve pain and function. (c) Office: Ongoing review and documentation of pain relief, functional status, appropriate medication use, and side effects. Pain assessment should include: current pain; the least reported pain over the period since last assessment; average pain; intensity of pain after taking the opioid; how long it takes for pain relief; and how long pain relief lasts. Satisfactory response to treatment may be indicated by the injured worker's decreased pain, increased level of function, or improved quality of life. Information from family members or other caregivers should be considered in determining the injured worker's response to treatment. The 4 A's for Ongoing Monitoring: Four domains have

been proposed as most relevant for ongoing monitoring of chronic pain injured workers on opioids: pain relief, side effects, physical and psychosocial functioning, and the occurrence of any potentially aberrant (or non-adherent) drug-related behaviors. These domains have been summarized as the "4 A's" (analgesia, activities of daily living, adverse side effects, and aberrant drug taking behaviors). The monitoring of these outcomes over time should affect therapeutic decisions and provide a framework for documentation of the clinical use of these controlled drugs. (Passik, 2000)

(d) Home: To aid in pain and functioning assessment, the injured worker should be requested to keep a pain diary that includes entries such as pain triggers, and incidence of end-of-dose pain. It should be emphasized that using this diary will help in tailoring the opioid dose. This should not be a requirement for pain management.

(e) Use of drug screening or injured worker treatment with issues of abuse, addiction, or poor pain control.

(f) Documentation of misuse of medications (doctor-shopping, uncontrolled drug escalation, drug diversion).

(g) Continuing review of overall situation with regard to non-opioid means of pain control.

(h) Consideration of a consultation with a multidisciplinary pain clinic if doses of opioids are required beyond what is usually required for the condition or pain does not improve on opioids in 3 months. Consider a psych consult if there is evidence of depression, anxiety or irritability.

Additionally, the MTUS states that continued use of opioids requires (a) the injured worker has returned to work, (b) the injured worker has improved functioning and pain. There is no current documentation of baseline pain, pain score with use of opioids, functional improvement on current regimen, side effects or review of potentially aberrant drug taking behaviors as outlined in the MTUS and as required for ongoing treatment. Therefore, at this time, the requirements for treatment have not been met and medical necessity has not been established.