

Case Number:	CM15-0026235		
Date Assigned:	02/18/2015	Date of Injury:	11/27/2012
Decision Date:	04/06/2015	UR Denial Date:	02/05/2015
Priority:	Standard	Application Received:	02/11/2015

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:
 State(s) of Licensure: North Carolina
 Certification(s)/Specialty: Family Practice

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The injured worker is a 40 year old female who sustained an industrial injury on 11/27/12. She reports back and leg pain. Treatments to date include medication, left knee surgery, and postoperative physical therapy to the knee. Diagnoses include persistent low back pain and left knee pain. In a progress note dated 01/07/15 the treating provider recommends continued medication including Norco, Naproxen, and Colace, a knee brace, and postoperative physical therapy, as well as follow up with the orthopedic surgeon. On 02/05/15 Utilization Review non-certified the physical therapy and Naproxen, citing MTUS guidelines.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

Eight (8) physical therapy: Upheld

Claims Administrator guideline: Decision based on MTUS Postsurgical Treatment Guidelines.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines physical medicine Page(s): 98-99.

Decision rationale: The California chronic pain medical treatment guidelines section on physical medicine states: Recommended as indicated below. Passive therapy (those treatment modalities that do not require energy expenditure on the part of the patient) can provide short term relief during the early phases of pain treatment and are directed at controlling symptoms such as pain ,inflammation and swelling and to improve the rate of healing soft tissue injuries. They can be used sparingly with active therapies to help control swelling, pain and inflammation during the rehabilitation process. Active therapy is based on the philosophy that therapeutic exercise and/or activity are beneficial for restoring flexibility, strength, endurance, function, range of motion, and can alleviate discomfort. Active therapy requires an internal effort by the individual to complete a specific exercise or task. This form of therapy may require supervision from a therapist or medical provider such as verbal, visual and/or tactile instruction(s). Patients are instructed and expected to continue active therapies at home as an extension of the treatment process in order to maintain improvement levels. Home exercise can include exercise with or without mechanical assistance or resistance and functional activities with assistive devices. (Colorado, 2002) (Airaksinen, 2006) Patient-specific hand therapy is very important in reducing swelling, decreasing pain, and improving range of motion in CRPS. (Li, 2005) The use of active treatment modalities (e.g., exercise, education, activity modification) instead of passive treatments is associated with substantially better clinical outcomes. In a large case series of patients with low back pain treated by physical therapists, those adhering to guidelines for active rather than passive treatments incurred fewer treatment visits, cost less, and had less pain and less disability. The overall success rates were 64.7% among those adhering to the active treatment recommendations versus 36.5% for passive treatment. (Fritz, 2007) Physical Medicine Guidelines allow for fading of treatment frequency (from up to 3 visits per week to 1 or less), plus active self-directed home Physical Medicine. Myalgia and myositis, unspecified (ICD9 729.1): 9-10 visits over 8 weeks. Neuralgia, neuritis, and radiculitis, unspecified (ICD9 729.2)8-10 visits over 4 weeks. Reflex sympathetic dystrophy (CRPS) (ICD9 337.2): 24 visits over 16 weeks. The requested amount of physical therapy is in excess of California chronic pain medical treatment guidelines. The patient has already completed a course of physical therapy. There is no explanation why the patient would need excess physical therapy and not be transitioned to active self-directed physical medicine. In the absence of such documentation, the request cannot be certified.

One (1) retrospective request (01/07/2015) prescription of Naproxen 550mg #30:
Overturned

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Naproxen (Naprosyn).

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines NSAIDs Page(s): 68.

Decision rationale: The California chronic pain medical treatment guideline section on NSAID therapy states: Recommended at the lowest dose for the shortest period in patients with moderate to severe pain. Acetaminophen may be considered for initial therapy for patients with mild to moderate pain, and in particular, for those with gastrointestinal, cardiovascular or renovascular risk factors. NSAIDs appear to be superior to acetaminophen, particularly for patients with

moderate to severe pain. There is no evidence to recommend one drug in this class over another based on efficacy. In particular, there appears to be no difference between traditional NSAIDs and COX-2 NSAIDs in terms of pain relief. The main concern of selection is based on adverse effects. COX-2 NSAIDs have fewer GI side effects at the risk of increased cardiovascular side effects, although the FDA has concluded that long-term clinical trials are best interpreted to suggest that cardiovascular risk occurs with all NSAIDs and is a class effect (with naproxyn being the safest drug). There is no evidence of long-term effectiveness for pain or function. (Chen, 2008) (Laine, 2008) This medication is recommended at the lowest possible dose for the shortest period of time. The duration of "shortest period of time" is not defined in the California MTUS. The patient has no mentioned cardiovascular, renovascular or gastrointestinal side-effects or risk factors. The dosage prescribed is within recommendations. Therefore, the request is certified.