

Case Number:	CM15-0029983		
Date Assigned:	02/24/2015	Date of Injury:	08/08/2006
Decision Date:	04/20/2015	UR Denial Date:	02/05/2015
Priority:	Standard	Application Received:	02/18/2015

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:
 State(s) of Licensure: New Jersey, Michigan, California
 Certification(s)/Specialty: Neurology, Neuromuscular Medicine

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The injured worker is a 49 year old male, who sustained an industrial injury on 8/08/2006. The mechanism of injury was not noted. reported. The injured worker was diagnosed as having cervical spinal stenosis and post cervical laminectomy syndrome. Treatment to date has included surgical and conservative treatments (cervical spine in 2009), including medications, physical therapy, and acupuncture. Currently, the injured worker complains of neck pain. Pain was rated 3/10 with medications and 4/10 without. He stated that he would like to decrease Percocet use to twice daily. He reported acupuncture sessions helping with pain. Current medications for pain included Percocet four times daily as needed and Orphenadrine ER. He continued to work full time. Exam of the cervical spine noted restricted range of motion and tenderness and tight muscle band on both sides. Sensory exam noted temperature sensation decrease over the C7-8 dermatome on the right. Exam of the bilateral shoulders noted positive Hawkin's and Neer tests.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

Percocet 10/325mg twice daily as needed, quantity 60 with one refill: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Opioid Use for Chronic pain.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Criteria for use of opioids Page(s): 179.

Decision rationale: According to MTUS guidelines, ongoing use of opioids should follow specific rules: (a) Prescriptions from a single practitioner taken as directed, and all prescriptions from a single pharmacy. (b) The lowest possible dose should be prescribed to improve pain and function. (c) Office: Ongoing review and documentation of pain relief, functional status, appropriate medication use, and side effects. Pain assessment should include: current pain; the least reported pain over the period since last assessment; average pain; intensity of pain after taking the opioid; how long it takes for pain relief; and how long pain relief lasts. Satisfactory response to treatment may be indicated by the patient's decreased pain, increased level of function, or improved quality of life. Information from family members or other caregivers should be considered in determining the patient's response to treatment. The 4 A's for Ongoing Monitoring: Four domains have been proposed as most relevant for ongoing monitoring of chronic pain patients on opioids: pain relief, side effects, physical and psychosocial functioning, and the occurrence of any potentially aberrant (or non adherent) drug-related behaviors. These domains have been summarized as the "4 A's" (analgesia, activities of daily living, adverse side effects, and aberrant drug taking behaviors). The monitoring of these outcomes over time should affect therapeutic decisions and provide a framework. The patient have been using opioids for long period of time without recent documentation of full control of pain and without any documentation of functional or quality of life improvement. There is no clear documentation of patient improvement in level of function, quality of life, adequate follow up for absence of side effects and aberrant behavior with a previous use of narcotics. There is no justification for the use of several narcotics. Therefore the prescription of Percocet 10/325mg, #60, with 1 refill is not medically necessary.