

Case Number:	CM15-0215194		
Date Assigned:	11/05/2015	Date of Injury:	08/26/2005
Decision Date:	12/21/2015	UR Denial Date:	10/13/2015
Priority:	Standard	Application Received:	11/02/2015

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:
 State(s) of Licensure: California, Oregon, Washington
 Certification(s)/Specialty: Orthopedic Surgery

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The injured worker is a 55-year-old male who sustained an industrial injury on August 26, 2005. Medical records indicated that the injured worker was treated for back and bilateral buttock pain. Medical diagnoses include degenerative disc disease, lumbar disc displacement. In the provider notes dated November 5, 2013 to November 6, 2013, the injured worker was admitted through the emergency room for management of back pain. The injured worker complained of a three-day history of gradual onset of severe, constant back pain radiating to both buttocks. He rates his pain 9 on the pain scale. He states his pain has not been controlled with oral narcotics. He states he is to have a future laminectomy. On exam, the documentation stated that there was soft tissue tenderness in the thoracic and lumbar area with limited range of motion in the back. He "is unable to ambulate to any significant distance and is unable to tolerate the pain on an outpatient basis." He is "diaphoretic and short of breath due to the severe pain." The treatment plan is for pain management and possible epidural injection. A Request for Authorization was submitted for retrospective 62310 inject spine CY 64483-inj foramen epidural lumbar. The Utilization Review dated October 13, 2015 denied the request for retrospective 62310 inject spine CY 64483-inj foramen epidural lumbar.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

Retrospective Injection spine C/T injection foramen epidural lumbar spine C/T (DOS 11/7/13): Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Medical Treatment 2009, Section(s): Epidural steroid injections (ESIs).

MAXIMUS guideline: Decision based on MTUS Chronic Pain Medical Treatment 2009, Section(s): Epidural steroid injections (ESIs).

Decision rationale: According to the CA MTUS Chronic Pain Medical Treatment Guidelines, Epidural injections are recommended as an option for treatment of radicular pain (defined as pain in dermatomal distribution with corroborative findings of radiculopathy). Specifically the guidelines state that radiculopathy must be documented by physical examination and corroborated by imaging studies and/or electrodiagnostic testing. Research has now shown that, on average, less than two injections are required for a successful ESI outcome. Current recommendations suggest a second epidural injection if partial success is produced with the first injection and a third ESI is rarely recommended. Epidural steroid injection can offer short-term pain relief and use should be in conjunction with other rehab efforts, including continuing a home exercise program. The American Academy of Neurology recently concluded that epidural steroid injections may lead to an improvement in radicular lumbosacral pain between 2 and 6 weeks following the injection, but they do not affect impairment of function or the need for surgery and do not provide long-term pain relief beyond 3 months. In addition, there must be demonstration of unresponsiveness to conservative treatment (exercises, physical methods, NSAIDs and muscle relaxants). The purpose of ESI is to reduce pain and inflammation, restoring range of motion and thereby facilitating progress in more active treatment programs, and avoiding surgery, but this treatment alone offers no significant long-term functional benefit. CA MTUS criteria for the use of Epidural steroid injections: 1) Radiculopathy must be documented by physical examination and corroborated by imaging studies and/or electrodiagnostic testing. 2) Initially unresponsive to conservative treatment (exercises, physical methods, NSAIDs and muscle relaxants). 3) Injections should be performed using fluoroscopy (live x-ray) for guidance. 4) If used for diagnostic purposes, a maximum of two injections should be performed. A second block is not recommended if there is inadequate response to the first block. Diagnostic blocks should be at an interval of at least one to two weeks between injections. 5) No more than two nerve root levels should be injected using transforaminal blocks. 6) No more than one interlaminar level should be injected at one session. 7) In the therapeutic phase, repeat blocks should be based on continued objective documented pain and functional improvement, including at least 50% pain relief with associated reduction of medication use for six to eight weeks, with a general recommendation of no more than 4 blocks per region per year. 8) Current research does not support a series-of-three injections in either the diagnostic or therapeutic phase. We recommend no more than 2 ESI injections. In this case the exam notes provided do not demonstrate a failure of conservative management nor a clear evidence of a dermatomal distribution of radiculopathy. Therefore, an epidural steroid injection is not medically necessary.