

Case Number:	CM15-0221314		
Date Assigned:	11/17/2015	Date of Injury:	08/19/2010
Decision Date:	12/31/2015	UR Denial Date:	10/29/2015
Priority:	Standard	Application Received:	11/11/2015

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:
 State(s) of Licensure: California, Oregon, Washington
 Certification(s)/Specialty: Orthopedic Surgery

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The injured worker is a 60 year old male, who sustained an industrial injury on 8-19-10. The injured worker was diagnosed as having bilateral ulnar neuropathy, cervical radiculopathy with weakness at right C6, and multilevel disc protrusion or herniation worse at C5-6. Treatment to date has included physical therapy, a home exercise program, and medication including Ambien, Norco, and Ibuprofen. The injured worker had been taking Ambien, Norco, and Ibuprofen since at least September 2014. On 8-28-15, the injured worker complained of neck pain with radiation to the right arm with numbness and tingling. On 10-23-15, the treating physician requested authorization for Norco 10-325mg #90, Ibuprofen 800mg #60, and Ambien 10mg #30. On 10-29-15 the requests were non-certified by utilization review.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

Norco 10/325 mg Qty 90: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Medical Treatment 2009, Section(s): Opioids (Classification).

MAXIMUS guideline: Decision based on MTUS Chronic Pain Medical Treatment 2009, Section(s): Opioids, criteria for use, Opioids, specific drug list. Decision based on Non-MTUS Citation Official Disability Guidelines (ODG) Pain / Opioids criteria for use.

Decision rationale: According to the CA MTUS/Chronic Pain Medical Treatment Guidelines, opioids (criteria for use & specific drug list): A therapeutic trial of opioids should not be employed until the patient has failed a trial of non-opioid analgesics. The patient should have at least one physical and psychosocial assessment by the treating doctor (and a possible second opinion by a specialist) to assess whether a trial of opioids should occur. Before initiating therapy, the patient should set goals, and the continued use of opioids should be contingent on meeting these goals. Guidelines recommend ongoing review and documentation of pain relief, functional status, appropriate medication use, and side effects. Pain assessment should include: current pain; the least reported pain over the period since last assessment; average pain; intensity of pain after taking the opioid; how long it takes for pain relief; and how long pain relief lasts. Satisfactory response to treatment may be indicated by the patient's decreased pain, increased level of function, or improved quality of life. The 4 A's for Ongoing Monitoring include analgesia, activities of daily living, adverse side effects, and aberrant drug-taking behaviors. Opioids may be continued if the patient has returned to work and the patient has improved function/pain. The ODG-TWC pain section comments specifically on criteria for the use of drug screening for ongoing opioid treatment. The ODG Pain / Opioids for chronic pain states "According to a major NIH systematic review, there is insufficient evidence to support the effectiveness of long-term opioid therapy for improving chronic pain, but emerging data support a dose-dependent risk for serious harms." ODG criteria (Pain / Opioids criteria for use) for continuing use of opioids include: "(a) If the patient has returned to work, (b) If the patient has improved functioning and pain." Based upon the records reviewed there is insufficient evidence to support the medical necessity of chronic narcotic use. There is lack of demonstrated functional improvement, percentage of relief, demonstration of urine toxicology compliance, return to work, or increase in activity from the exam note of 8/28/15. Therefore, the request is not medically necessary.

Ibuprofen 800 mg Qty 60: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Medical Treatment 2009, Section(s): NSAIDs (non-steroidal anti-inflammatory drugs).

MAXIMUS guideline: Decision based on MTUS Chronic Pain Medical Treatment 2009, Section(s): NSAIDs (non-steroidal anti-inflammatory drugs).

Decision rationale: According to the CA/MTUS Chronic Pain Medical Treatment Guidelines, page 67, NSAIDs, specific recommendations are for "Osteoarthritis (including knee and hip): Recommended at the lowest dose for the shortest period in patients with moderate to severe pain. Acetaminophen may be considered for initial therapy for patients with mild to moderate pain, and in particular, for those with gastrointestinal, cardiovascular or renovascular risk factors. NSAIDs appear to be superior to acetaminophen, particularly for patients with moderate to severe pain. There is no evidence to recommend one drug in this class over another based on efficacy. In particular, there appears to be no difference between traditional NSAIDs and COX-2

NSAIDs in terms of pain relief. The main concern of selection is based on adverse effects. COX- 2 NSAIDs have fewer GI side effects at the risk of increased cardiovascular side effects, although the FDA has concluded that long-term clinical trials are best interpreted to suggest that cardiovascular risk occurs with all NSAIDs and is a class effect (with naproxyn being the safest drug). There is no evidence of long-term effectiveness for pain or function. (Chen, 2008) (Laine, 2008)" In this case after review of the medical records from 8/28/15 there is insufficient evidence to support functional improvement on Ibuprofen or osteoarthritis to warrant usage. Therefore the the request is not medically necessary.

Ambien 10 mg Qty 30: Upheld

Claims Administrator guideline: The Claims Administrator did not base their decision on the MTUS. Decision based on Non-MTUS Citation Official Disability Guidelines: Pain (Chronic) - Zolpidem (Ambien).

MAXIMUS guideline: The Expert Reviewer did not base their decision on the MTUS. Decision based on Non-MTUS Citation Official Disability Guidelines (ODG) Pain Section, Zolpidem (Ambien).

Decision rationale: CA MTUS/ACOEM is silent on the issue of Ambien. According to the ODG, Pain Section, Zolpidem (Ambien) is a prescription short-acting non-benzodiazepine hypnotic, which is approved for the short-term (usually two to six weeks) treatment of insomnia. Proper sleep hygiene is critical to the individual with chronic pain and often is hard to obtain. Various medications may provide short-term benefit. While sleeping pills, so-called minor tranquilizers, and anti-anxiety agents are commonly prescribed in chronic pain, pain specialists rarely, if ever, recommend them for long-term use. They can be habit-forming, and they may impair function and memory more than opioid pain relievers. There is also concern that they may increase pain and depression over the long-term. There is no evidence in the records from 8/28/15 of insomnia to warrant Ambien. ODG guidelines do not recommend long-term use of Ambien. Therefore, the request is not medically necessary.