

Case Number:	CM15-0222796		
Date Assigned:	11/18/2015	Date of Injury:	02/28/2012
Decision Date:	12/30/2015	UR Denial Date:	10/15/2015
Priority:	Standard	Application Received:	11/12/2015

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:

State(s) of Licensure: Massachusetts

Certification(s)/Specialty: Anesthesiology, Pain Management

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

This injured worker is a 51 year old female, who sustained an industrial injury on 02-28-2012. The injured worker was diagnosed as having left shoulder impingement syndrome. On medical records dated 10-08-2015, the subjective complaints were noted as left shoulder pain, numbness and tingling, radiating to neck associated with repetitive movement. Objective findings were noted as left shoulder with range of motion was decreased. Tenderness to palpation of the anterior shoulder, glenohumeral joint and posterior shoulder. There was muscle spasm of the anterior shoulder and posterior shoulder as well as a positive for impingement. Treatment to date included medication. Current medications were listed as Tramadol, Diclofenac Sodium and Pantoprazole. The Utilization Review (UR) was dated 10-15-2015. A Request for Authorization was submitted. The UR submitted for this medical review indicated that the request for pantoprazole 20 mg 1 BID #60, compound HS AGB-Amitriptyline 10%, Gabapentin 10%, Bupivacaine 5% in cream base #240 gms and compound FBD-Flurbiprofen 20%, Baclofen 10%, Dexamethasone 1%, Panthenol 0.5% in cream base #240 gms was non-certified.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

Pantoprazole 20 mg 1 BID #60: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Medical Treatment 2009, Section(s): NSAIDs, GI symptoms & cardiovascular risk.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Medical Treatment 2009, Section(s): NSAIDs, GI symptoms & cardiovascular risk.

Decision rationale: The MTUS makes the following recommendations for the use of proton pump inhibitors. Clinicians should weight the indications for NSAIDs against both GI and cardiovascular risk factors. Determine if the injured worker is at risk for gastrointestinal events: (1) age > 65 years; (2) history of peptic ulcer, GI bleeding or perforation; (3) concurrent use of ASA, corticosteroids, and/or an anticoagulant; or (4) high dose/multiple NSAID (e.g., NSAID + low-dose ASA). Recent studies tend to show that H. Pylori does not act synergistically with NSAIDS to develop gastroduodenal lesions. Recommendations Injured workers with no risk factor and no cardiovascular disease: Non-selective NSAIDs OK (e.g, ibuprofen, naproxen, etc.) Injured workers at intermediate risk for gastrointestinal events and no cardiovascular disease: (1) A non-selective NSAID with either a PPI (Proton Pump Inhibitor, for example, 20 mg omeprazole daily) or misoprostol (200 ug four times daily) or (2) a Cox-2 selective agent. Long-term PPI use (> 1 year) has been shown to increase the risk of hip fracture (adjusted odds ratio 1.44). Injured workers at high risk for gastrointestinal events with no cardiovascular disease: A Cox-2 selective agent plus a PPI if absolutely necessary. Injured workers at high risk of gastrointestinal events with cardiovascular disease: If GI risk is high the suggestion is for a low-dose Cox-2 plus low dose Aspirin (for cardio-protection) and a PPI. If cardiovascular risk is greater than GI risk the suggestion is naproxyn plus low-dose aspirin plus a PPI. Cardiovascular disease: A non-pharmacological choice should be the first option in injured workers with cardiac risk factors. It is then suggested that acetaminophen or aspirin be used for short term needs. An opioid also remains a short-term alternative for analgesia. Major risk factors (recent MI, or coronary artery surgery, including recent stent placement): If NSAID therapy is necessary, the suggested treatment is naproxyn plus low-dose aspirin plus a PPI. Mild to moderate risk factors: If long-term or high-dose therapy is required, full-dose naproxen (500 mg twice a day) appears to be the preferred choice of NSAID. If naproxyn is ineffective, the suggested treatment is (1) the addition of aspirin to naproxyn plus a PPI, or (2) a low-dose Cox-2 plus ASA. According to the records available for review the injured worker does not meet any of the guidelines required for the use of this medication therefore, at this time, the requirements for treatment have not been met and medical necessity has not been established.

Compound HS AGB-Amitriptyline 10%, Gabapentin 10%, Bupivacaine 5% in cream base #240 gms: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Medical Treatment 2009, Section(s): Topical Analgesics.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Medical Treatment 2009, Section(s): Topical Analgesics.

Decision rationale: According to the MTUS, there is little to no research to support the use of topical compounded creams. The use of these compounded agents requires knowledge of the specific analgesic effect of each agent and how it will be useful for the specific therapeutic goal required. Topical analgesics are largely experimental and there are a few randomized controlled trials to determine efficacy or safety. Therefore, at this time, the requirements for treatment have not been met and medical necessity has not been established.

Compound FBD-Flurbiprofen 20%, Baclofen 10%, Dexamethasone 1%, Panthenol 0.5% in cream base #240 gms: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Medical Treatment 2009, Section(s): Topical Analgesics.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Medical Treatment 2009, Section(s): Topical Analgesics.

Decision rationale: According to the MTUS, there is little to no research to support the use of topical compounded creams. The use of these compounded agents requires knowledge of the specific analgesic effect of each agent and how it will be useful for the specific therapeutic goal required. Topical analgesics are largely experimental and there are a few randomized controlled trials to determine efficacy or safety. Therefore, at this time, the requirements for treatment have not been met and medical necessity has not been established.