

Case Number:	CM15-0030225		
Date Assigned:	02/23/2015	Date of Injury:	03/20/2006
Decision Date:	04/07/2015	UR Denial Date:	02/04/2015
Priority:	Standard	Application Received:	02/18/2015

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:
 State(s) of Licensure: New Jersey, Michigan, California
 Certification(s)/Specialty: Neurology, Neuromuscular Medicine

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The injured worker is a 56 year old male, who sustained an industrial injury on 03/20/2006. The diagnoses have included post-laminectomy syndrome of the lumbar spine, lumbar radiculopathy, sacroiliitis, degeneration of the lumbar disks, and shoulder pain. Noted treatments to date have included surgeries, physical therapy, acupuncture, steroid injections, pool therapy, psychotherapy, and medications. No MRI report noted in received medical records. In a progress note dated 12/19/2014, the injured worker presented for his monthly medication management appointment. The treating physician reported the injured worker presented with complaints of chronic right jaw pain since receiving a recent injection along with lower back pain with radiation of pain down his right lower extremity. Utilization Review determination on 02/04/2015 modified the request for Monthly Medication Management Visits x 12 to 3 Monthly Medication Management Visits and non-certified the request for and Lumbar Trigger Point Injection with Ultrasound citing Medical Treatment Utilization Schedule Chronic Pain Medical Treatment Guidelines and Official Disability Guidelines.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

Monthly medication management visits x 12: Upheld

Claims Administrator guideline: The Claims Administrator did not base their decision on the MTUS. Decision based on Non-MTUS Citation Official Disability Guidelines.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Early intervention Page(s): 32-33.

Decision rationale: According to MTUS guidelines, the presence of red flags may indicate the need for specialty consultation. In addition, the requesting physician should provide a documentation supporting the medical necessity for a pain management evaluation with a specialist. The documentation should include the reasons, the specific goals and end point for using the expertise of a specialist. There is no clear documentation that the patient needs a monthly visit for medications management per MTUS criteria. There is no clear documentation that the patient had delayed recovery or a medical condition that require a monthly adjustment of the patient medications. The provider did not document the reasons, the specific goals and end point for follow up visits. Therefore, the request is not medically necessary.

Lumbar trigger point injections with ultrasound: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Trigger point injections.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Trigger point injections Page(s): 122.

Decision rationale: According to MTUS guidelines, trigger point injection is recommended only for myofascial pain syndrome as indicated below, with limited lasting value. Not recommended for radicular pain. Trigger point injections with an anesthetic such as bupivacaine are recommended for non-resolving trigger points, but the addition of a corticosteroid is not generally recommended. Not recommended for radicular pain. A trigger point is a discrete focal tenderness located in a palpable taut band of skeletal muscle, which produces a local twitch in response to stimulus to the band. Trigger points may be present in up to 33-50% of the adult population. Myofascial pain syndrome is a regional painful muscle condition with a direct relationship between a specific trigger point and its associated pain region. These injections may occasionally be necessary to maintain function in those with myofascial problems when myofascial trigger points are present on examination. Not recommended for typical back pain or neck pain. (Graff-Radford, 2004) (Nelemans-Cochrane, 2002) For fibromyalgia syndrome, trigger point injections have not been proven effective. (Goldenberg, 2004) Trigger point injections with a local anesthetic may be recommended for the treatment of chronic low back or neck pain with myofascial pain syndrome when all of the following criteria are met: (1) Documentation of circumscribed trigger points with evidence upon palpation of a twitch response as well as referred pain; (2) Symptoms have persisted for more than three months; (3) Medical management therapies such as ongoing stretching exercises, physical therapy, NSAIDs and muscle relaxants have failed to control pain; (4) Radiculopathy is not present (by exam, imaging, or neuro-testing); (5) Not more than 3-4 injections per session; (6) No repeat injections unless a greater than 50% pain relief is obtained for six weeks after an injection and there is documented evidence of functional improvement; (7) Frequency should not be at an

interval less than two months; (8) Trigger point injections with any substance (e.g., saline or glucose) other than local anesthetic with or without steroid are not recommended. There is no clear evidence of myofascial pain and trigger points over the lumbar and sciatic notch. There is no documentation of failure of oral medications or physical therapy in this case. Therefore, the request for lumbar Trigger point injection is not medically necessary.