MAXIMUS Federal Services

Case Number:	CM15-0033759		
Date Assigned:	02/27/2015	Date of Injury:	08/03/2002
Decision Date:	04/06/2015	UR Denial Date:	01/30/2015
Priority:	Standard	Application Received:	02/23/2015

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:
State(s) of Licensure: North Carolina
Certification(s)/Specialty: Family Practice

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The injured worker is a 50year old female, who sustained a work/ industrial injury on 8/3/02 while riding a motorcycle in a funeral procession when she was hit by a car and thrown to the ground and sustained numerous injuries to include, closed head injury, left tibia and fibula fracture, and right lower leg laceration. She has reported symptoms of right shoulder, neck, left knee, and ankle pain. Medical history includes gastrointestinal distress with weight loss and constipation and chronic anemia. Surgery included left tibia and fibula open reduction internal fixation (ORIF).The diagnoses have included partial thickness rotator cuff tear. Treatments to date included medication, physical therapy, conservative management, orthopedic consultation, and psychotherapy. Diagnostics included an Magnetic Resonance Imaging (MRI) from 11/24/09 noted no evidence of a full thickness rotator cuff tear with type 2 acromial morphology, no glenohumeral degenerative joint disease, minimal to mild degenerative joint disease of the acromioclavicular joint, no loose bodies, no labral tears, intact bicep tendon, intact subscapularis tendon, with an increased signal in the supraspinatus tendon. Computed Tomography (CT) of the left knee reported anterior tibial metaphysic with healed fracture. Medications included Norco, Prilosec, Zanaflex, Neurontin, and Prozac. Examination noted full range of motion with a 20% loss of abduction and forward flexion, mild loss of internal rotation, neurological exam was intact, on the right upper extremity, Neer, Hawkin's tests were positive, cross arm test, Speed test, Yergason, O'Brien tests were negative. On 1/30/15, Utilization Review modified Norco 10/325mg #120 to Norco 10/325mg #48, citing the California Medical treatment Utilization Schedule (MTUS) Guidelines.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

Norco 10/325 MG #120: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Opioids.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines opioids Page(s): 76-84.

Decision rationale: The California chronic pain medical treatment guidelines section on opioids states for ongoing management: On-Going Management. Actions Should Include: (a) Prescriptions from a single practitioner taken as directed, and all prescriptions from a single pharmacy. (b) The lowest possible dose should be prescribed to improve pain and function. (c) Office: Ongoing review and documentation of pain relief, functional status, appropriate medication use, and side effects. Pain assessment should include: current pain; the least reported pain over the period since last assessment; average pain; intensity of pain after taking the opioid; how long it takes for pain relief; and how long pain relief lasts. Satisfactory response to treatment may be indicated by the patient's decreased pain, increased level of function, or improved quality of life. Information from family members or other caregivers should be considered in determining the patient's response to treatment. The 4 A's for Ongoing Monitoring: Four domains have been proposed as most relevant for ongoing monitoring of chronic pain patients on opioids: pain relief, side effects, physical and psychosocial functioning, and the occurrence of any potentially aberrant (or non-adherent) drug-related behaviors. These domains have been summarized as the "4 A's" (analgesia, activities of daily living, adverse side effects, and aberrant drug taking behaviors). The monitoring of these outcomes over time should affect therapeutic decisions and provide a framework for documentation of the clinical use of these controlled drugs. (Passik, 2000) (d) Home: To aid in pain and functioning assessment, the patient should be requested to keep a pain dairy that includes entries such as pain triggers, and incidence of end-of-dose pain. It should be emphasized that using this diary will help in tailoring the opioid dose. This should not be a requirement for pain management. (e) Use of drug screening or inpatient treatment with issues of abuse, addiction, or poor pain control. (f) Documentation of misuse of medications (doctor- shopping, uncontrolled drug escalation, drug diversion). (g) Continuing review of overall situation with regard to non-opioid means of pain control. (h) Consideration of a consultation with a multidisciplinary pain clinic if doses of opioids are required beyond what is usually required for the condition or pain does not improve on opioids in 3 months. Consider a psych consult if there is evidence of depression, anxiety or irritability. Consider an addiction medicine consult if there is evidence of substance misuse. When to Continue Opioids (a) If the patient has returned to work (b) If the patient has improved functioning and pain(Washington, 2002) (Colorado, 2002) (Ontario, 2000) (VA/DoD, 2003) (Maddox-AAPM/APS, 1997) (Wisconsin, 2004) (Warfield, 2004)The long-term use of this medication class is not recommended per the California MTUS unless there documented evidence of benefit with measurable outcome measures and improvement in function. There is a documented improvement in subjective VAS scores on the medication There however are no objective measurements of improvement in function. Therefore criteria for the ongoing use of opioids have not been met and the request is not certified.