

Case Number:	CM15-0033873		
Date Assigned:	02/27/2015	Date of Injury:	09/26/2011
Decision Date:	04/08/2015	UR Denial Date:	02/19/2015
Priority:	Standard	Application Received:	02/23/2015

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:

State(s) of Licensure: Massachusetts

Certification(s)/Specialty: Anesthesiology, Pain Management

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The injured worker is a year old female who sustained a work related injury on September 26, 2011, incurring left upper extremity pain and right upper extremity pain after lifting and repetitive motions. Treatments included splinting, physical therapy, nerve blocks, home exercise program, icing and stretching and pain and nerve medications. She was diagnosed with carpal tunnel syndrome and complex regional pain syndrome in the left hand. The injured worker underwent carpal tunnel release. Currently, in December, 2014, the injured worker complained of ongoing pain in the left upper extremity, numbness, and tingling. On February 27, 2015, a request for one prescription for Omeprazole 20 mg, #30 with 4 refills was modified to one prescription of Omeprazole 20 mg, #30, without refills by Utilization Review, noting the California Medical Treatment Utilization Schedule Guidelines Chronic Pain Medical Treatment Guidelines.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

Omeprazole 10mg, QTY: 30 with 4 refills: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines NSAIDs, GI symptoms & cardiovascular risk.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines PPI
Page(s): 68-69.

Decision rationale: PPI, p68-69 The MTUS makes the following recommendations for the use of proton pump inhibitors. Clinicians should weight the indications for NSAIDs against both GI and cardiovascular risk factors. Determine if the injured worker is at risk for gastrointestinal events: (1) age > 65 years; (2) history of peptic ulcer, GI bleeding or perforation; (3) concurrent use of ASA, corticosteroids, and/or an anticoagulant; or (4) high dose/multiple NSAID (e.g., NSAID + low-dose ASA). Recent studies tend to show that H. Pylori does not act synergistically with NSAIDs to develop gastroduodenal lesions. Recommendations Injured workers with no risk factor and no cardiovascular disease: Non-selective NSAIDs OK (e.g, ibuprofen, naproxen, etc.) Injured workers at intermediate risk for gastrointestinal events and no cardiovascular disease:(1) A non-selective NSAID with either a PPI (Proton Pump Inhibitor, for example, 20 mg omeprazole daily) or misoprostol (200 g four times daily) or (2) a Cox-2 selective agent. Long-term PPI use (> 1 year) has been shown to increase the risk of hip fracture (adjusted odds ratio 1.44). Injured workers at high risk for gastrointestinal events with no cardiovascular disease: A Cox-2 selective agent plus a PPI if absolutely necessary. Injured workers at high risk of gastrointestinal events with cardiovascular disease: If GI risk is high the suggestion is for a low-dose Cox-2 plus low dose Aspirin (for cardioprotection) and a PPI. If cardiovascular risk is greater than GI risk the suggestion is naproxyn plus low-dose aspirin plus a PPI. Cardiovascular disease: A non-pharmacological choice should be the first option in injured workers with cardiac risk factors. It is then suggested that acetaminophen or aspirin be used for short term needs. An opioid also remains a short-term alternative for analgesia. Major risk factors (recent MI, or coronary artery surgery, including recent stent placement): If NSAID therapy is necessary, the suggested treatment is naproxyn plus low-dose aspirin plus a PPI. Mild to moderate risk factors: If long-term or high-dose therapy is required, full-dose naproxen (500 mg twice a day) appears to be the preferred choice of NSAID. If naproxyn is ineffective, the suggested treatment is (1) the addition of aspirin to naproxyn plus a PPI, or (2) a low-dose Cox-2 plus ASA. According to the records available for review the injured worker does not meet any of the guidelines required for the use of this medication therefore, at this time, the requirements for treatment have not been met and medical necessity has not been established.