

Case Number:	CM15-0035760		
Date Assigned:	03/04/2015	Date of Injury:	11/29/2009
Decision Date:	04/14/2015	UR Denial Date:	02/02/2015
Priority:	Standard	Application Received:	02/25/2015

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:
 State(s) of Licensure: New Jersey, Michigan, California
 Certification(s)/Specialty: Neurology, Neuromuscular Medicine

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The injured worker is a 58 year old female, who sustained an industrial injury on 11/29/09. She has reported neck, low back, shoulders and wrists injuries. The diagnoses have included cervical strain/sprain, lumbosacral/thoracic radiculitis, lumbar strain, bilateral shoulder and wrist tendinosis and cervical radiculopathy. Treatment to date has included medications, cervical Epidural Steroid Injection (ESI) and Home Exercise Program (HEP). Currently, as per physician progress note dated 1/9/15, the injured worker complains of continued pain in the neck, low back, shoulders and bilateral wrists. She was having difficulty doing much activity above shoulder level or with her upper extremities. The current medications include Hydrocodone and Prilosec. Physical exam revealed spasm and tenderness of the lumbar spine with decreased range of motion. Straight leg raise causes back pain. Tenderness over the bilateral wrists was noted with Tinel's and Phalen's tests which were positive. She is not currently undergoing any therapy or other modes of treatment. There were no recent diagnostics or physical therapy sessions noted. Work status was temporary total disabled. On 2/2/15 Utilization Review non-certified a request for Norco 10/325mg #60 and Prilosec 20mg #30, noting the (MTUS) Medical Treatment Utilization Schedule chronic pain guidelines were cited.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

Norco 10/325mg #60: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Hydrocodone/Acetaminophen, Opioids Page(s): 76-79.

Decision rationale: According to MTUS guidelines, Norco (Hydrocodone/Acetaminophen) is a synthetic opioid indicated for the pain management but not recommended as a first line oral analgesic. In addition and according to MTUS guidelines, ongoing use of opioids should follow specific rules: (a) Prescriptions from a single practitioner taken as directed, and all prescriptions from a single pharmacy. (b) The lowest possible dose should be prescribed to improve pain and function. (c) Office: Ongoing review and documentation of pain relief, functional status, appropriate medication use, and side effects. Four domains have been proposed as most relevant for ongoing monitoring of chronic pain patients on opioids: pain relief, side effects, physical and psychosocial functioning, and the occurrence of any potentially aberrant (or non adherent) drug-related behaviors. These domains have been summarized as the "4 A's" (analgesia, activities of daily living, adverse side effects, and aberrant drug taking behaviors). The monitoring of these outcomes over time should affect therapeutic decisions and provide a framework. According to the patient file, there is no objective documentation of pain and functional improvement to justify continuous use of Norco. Norco was used for longtime without documentation of functional improvement or evidence of return to work or improvement of activity of daily living. Therefore, the prescription of Norco 10/325mg #60 is not medically necessary.

Prilosec 20mg #30: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines NSAIDs, GI symptoms & cardiovascular risk Page(s): 68.

Decision rationale: According to MTUS guidelines, Omeprazole is indicated when NSAID are used in patients with intermediate or high risk for gastrointestinal events. The risk for gastrointestinal events are: (1) age > 65 years; (2) history of peptic ulcer, GI bleeding or perforation; (3) concurrent use of ASA, corticosteroids, and/or an anticoagulant; or (4) high dose/multiple NSAID (e.g., NSAID + low-dose ASA). Recent studies tend to show that H. Pylori does not act synergistically with NSAIDS to develop gastroduodenal lesions. There is no documentation that the patient have GI issue that requires the use of prilosec. There is no documentation in the patient's chart supporting that she is at intermediate or high risk for developing gastrointestinal events. Therefore, the request for Prilosec 20mg #30 prescription is not medically necessary.

