

Case Number:	CM15-0035814		
Date Assigned:	03/04/2015	Date of Injury:	06/21/2007
Decision Date:	04/13/2015	UR Denial Date:	02/03/2015
Priority:	Standard	Application Received:	02/25/2015

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:
 State(s) of Licensure: New Jersey, Michigan, California
 Certification(s)/Specialty: Neurology, Neuromuscular Medicine

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The injured worker is a 55 year old male, who sustained an industrial injury on June 21, 2007. The injured worker had sustained a neck and bilateral shoulder injury. The diagnoses have included severe degenerative disc disease of the lumbar three-lumbar four levels and status post cervical discectomy. Treatment to date has included medications, imaging studies and physical therapy. Current documentation dated November 25, 2014 notes that the injured worker reported neck and bilateral shoulder pain. The neck pain was described as a continuous numbness radiating to the chest and head. The injured worker noted that his symptoms had improved from the prior visit secondary to cervical spine surgery. The bilateral shoulder pain was described as occasional, stabbing and rated a two out of ten on the Visual Analogue Scale. The pain was noted to radiate to the neck. Physical examination was unchanged from the prior visit. On February 3, 2015 Utilization Review non-certified a request for physical therapy two times a week for six weeks to the cervical spine and physical therapy two times a week for six weeks to the lumbar spine. The Official Disability Guidelines were cited.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

Physical therapy 2x6 cervical: Upheld

Claims Administrator guideline: The Claims Administrator did not cite any medical evidence for its decision.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Physical Medicine Page(s): 98.

Decision rationale: According to MTUS guidelines, Physical Medicine is "Recommended as indicated below. Passive therapy (those treatment modalities that do not require energy expenditure on the part of the patient) can provide short term relief during the early phases of pain treatment and are directed at controlling symptoms such as pain, inflammation and swelling and to improve the rate of healing soft tissue injuries. They can be used sparingly with active therapies to help control swelling, pain and inflammation during the rehabilitation process. Active therapy is based on the philosophy that therapeutic exercise and/or activity are beneficial for restoring flexibility, strength, endurance, function, range of motion, and can alleviate discomfort. Active therapy requires an internal effort by the individual to complete a specific exercise or task. This form of therapy may require supervision from a therapist or medical provider such as verbal, visual and/or tactile instruction(s). Patients are instructed and expected to continue active therapies at home as an extension of the treatment process in order to maintain improvement levels. Home exercise can include exercise with or without mechanical assistance or resistance and functional activities with assistive devices. (Colorado, 2002) (Airaksinen, 2006) Patient-specific hand therapy is very important in reducing swelling, decreasing pain, and improving range of motion in CRPS. (Li, 2005) The use of active treatment modalities (e.g., exercise, education, activity modification) instead of passive treatments is associated with substantially better clinical outcomes. In a large case series of patients with low back pain treated by physical therapists, those adhering to guidelines for active rather than passive treatments incurred fewer treatment visits, cost less, and had less pain and less disability. The overall success rates were 64.7% among those adhering to the active treatment recommendations versus 36.5% for passive treatment (Fritz, 2007)." There is no documentation of the efficacy and outcome of previous physical therapy sessions. There is no documentation that the patient cannot perform home exercise. Therefore, the request for Physical therapy 2x6 cervical is not medically necessary.

Physical therapy 2x6 lumbar spine: Upheld

Claims Administrator guideline: The Claims Administrator did not cite any medical evidence for its decision.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Physical Medicine Page(s): 98.

Decision rationale: According to MTUS guidelines, Physical Medicine is "Recommended as indicated below. Passive therapy (those treatment modalities that do not require energy expenditure on the part of the patient) can provide short term relief during the early phases of pain treatment and are directed at controlling symptoms such as pain, inflammation and swelling and to improve the rate of healing soft tissue injuries. They can be used sparingly with active therapies to help control swelling, pain and inflammation during the rehabilitation process.

Active therapy is based on the philosophy that therapeutic exercise and/or activity are beneficial for restoring flexibility, strength, endurance, function, range of motion, and can alleviate discomfort. Active therapy requires an internal effort by the individual to complete a specific exercise or task. This form of therapy may require supervision from a therapist or medical provider such as verbal, visual and/or tactile instruction(s). Patients are instructed and expected to continue active therapies at home as an extension of the treatment process in order to maintain improvement levels. Home exercise can include exercise with or without mechanical assistance or resistance and functional activities with assistive devices. (Colorado, 2002) (Airaksinen, 2006) Patient-specific hand therapy is very important in reducing swelling, decreasing pain, and improving range of motion in CRPS. (Li, 2005) The use of active treatment modalities (e.g., exercise, education, activity modification) instead of passive treatments is associated with substantially better clinical outcomes. In a large case series of patients with low back pain treated by physical therapists, those adhering to guidelines for active rather than passive treatments incurred fewer treatment visits, cost less, and had less pain and less disability. The overall success rates were 64.7% among those adhering to the active treatment recommendations versus 36.5% for passive treatment (Fritz, 2007)." There is no documentation of pain and functional improvement with previous physical therapy sessions. There is no documentation that the patient cannot perform home exercise. Therefore, the request for Physical therapy 2x6 lumbar spine is not medically necessary.