

Case Number:	CM15-0038829		
Date Assigned:	03/09/2015	Date of Injury:	04/15/2013
Decision Date:	04/10/2015	UR Denial Date:	02/12/2015
Priority:	Standard	Application Received:	03/02/2015

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:
 State(s) of Licensure: North Carolina
 Certification(s)/Specialty: Family Practice

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

This 52-year-old woman sustained an industrial injury on 4/15/2013. The mechanism of injury is not detailed. Current diagnoses include lateral meniscus tear, esophagus reflux, and contusion of knee. Treatment has included oral medications and use of a cane. Physician notes on a PR-2 dated 1/28/2014 show complaints of moderate low back pain. Recommendations include Omeprazole, Norco, modified work duties, and post-operative physical therapy to begin two weeks after surgical intervention. On 2/12/2015, Utilization Review evaluated prescriptions for four sessions of psychotherapy, and an additional 12 sessions of physical therapy for the left knee, that were submitted on 2/24/2015. The MTUS, ACOEM Guidelines, (or ODG) was cited. The requests were denied and subsequently appealed to Independent Medical Review.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

Psychotherapy; four (4) sessions: Overturned

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines psychological treatment Page(s): 101-102.

Decision rationale: The California chronic pain medical treatment guidelines section on psychological treatment states: Recommended for appropriately identified patients during treatment for chronic pain. Psychological intervention for chronic pain includes setting goals, determining appropriateness of treatment, conceptualizing a patient's pain beliefs and coping styles, assessing psychological and cognitive function, and addressing co-morbid mood disorders (such as depression, anxiety, panic disorder, and posttraumatic stress disorder). Cognitive behavioral therapy and self-regulatory treatments have been found to be particularly effective. Psychological treatment incorporated into pain treatment has been found to have a positive short-term effect on pain interference and long-term effect on return to work. The following stepped-care approach to pain management that involves psychological intervention has been suggested: Step 1: Identify and address specific concerns about pain and enhance interventions that emphasize self-management. The role of the psychologist at this point includes education and training of pain care providers in how to screen for patients that may need early psychological intervention. Step 2: Identify patients who continue to experience pain and disability after the usual time of recovery. At this point a consultation with a psychologist allows for screening, assessment of goals, and further treatment options, including brief individual or group therapy. Step 3: Pain is sustained in spite of continued therapy (including the above psychological care). Intensive care may be required from mental health professions allowing for a multidisciplinary treatment approach. See also Multi-disciplinary pain programs. See also ODG Cognitive Behavioral Therapy (CBT) Guidelines. (Otis, 2006) (Townsend, 2006) (Kerns, 2005) (Flor, 1992) (Morley, 1999) (Ostelo, 2005) Psychological treatment in particular cognitive behavioral therapy has been found to be particularly effective in the treatment of chronic pain. As this patient has continued ongoing pain, this service is indicated per the California MTUS and thus is certified.

Physical therapy; additional twelve (12) sessions, left knee: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Physical medicine.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines physical medicine Page(s): 98-99.

Decision rationale: The California chronic pain medical treatment guidelines section on physical medicine states: Recommended as indicated below. Passive therapy (those treatment modalities that do not require energy expenditure on the part of the patient) can provide short term relief during the early phases of pain treatment and are directed at controlling symptoms such as pain, inflammation and swelling and to improve the rate of healing soft tissue injuries. They can be used sparingly with active therapies to help control swelling, pain and inflammation during the rehabilitation process. Active therapy is based on the philosophy that therapeutic exercise and/or activity are beneficial for restoring flexibility, strength, endurance, function, range of motion, and can alleviate discomfort. Active therapy requires an internal effort by the individual to complete a specific exercise or task. This form of therapy may require supervision

from a therapist or medical provider such as verbal, visual and/or tactile instruction(s). Patients are instructed and expected to continue active therapies at home as an extension of the treatment process in order to maintain improvement levels. Home exercise can include exercise with or without mechanical assistance or resistance and functional activities with assistive devices. (Colorado, 2002) (Airaksinen, 2006) Patient-specific hand therapy is very important in reducing swelling, decreasing pain, and improving range of motion in CRPS. (Li, 2005) The use of active treatment modalities (e.g., exercise, education, activity modification) instead of passive treatments is associated with substantially better clinical outcomes. In a large case series of patients with low back pain treated by physical therapists, those adhering to guidelines for active rather than passive treatments incurred fewer treatment visits, cost less, and had less pain and less disability. The overall success rates were 64.7% among those adhering to the active treatment recommendations versus 36.5% for passive treatment. (Fritz, 2007) Physical Medicine Guidelines: Allow for fading of treatment frequency (from up to 3 visits per week to 1 or less), plus active self-directed home Physical Medicine. Myalgia and myositis, unspecified (ICD9 729.1): 9-10 visits over 8 weeks. Neuralgia, neuritis, and radiculitis, unspecified (ICD9 729.2) 8-10 visits over 4 weeks. Reflex sympathetic dystrophy (CRPS) (ICD9 337.2): 24 visits over 16 weeks. The requested amount of physical therapy is in excess of California chronic pain medical treatment guidelines. There is no explanation why the patient would need excess physical therapy and not be transitioned to active self-directed physical medicine. In the absence of such documentation, the request cannot be certified.