

| | | | |
|-----------------------|--------------|------------------------------|------------|
| Case Number: | CM15-0042098 | | |
| Date Assigned: | 03/12/2015 | Date of Injury: | 05/25/2000 |
| Decision Date: | 04/20/2015 | UR Denial Date: | 02/09/2015 |
| Priority: | Standard | Application Received: | 03/05/2015 |

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:

State(s) of Licensure: Indiana

Certification(s)/Specialty: Preventive Medicine, Occupational Medicine

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The injured worker is a 53-year-old male, who sustained an industrial injury on 5/25/2000. She reported a low back injury. The mechanism of injury was not provided for review. The injured worker was diagnosed as having chronic pain syndrome, lumbosacral neuritis or radiculitis, lumbosacral degeneration, lumbago, myalgia and myositis, dyesthesias and muscle spasm. Treatment to date has included physical therapy, epidural steroid injection and medication management. Currently, a progress note from the treating provider dated 2/3/2015 indicates the injured worker reported low back pain with left lower extremity pain.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

Lorazepam 1 mg #60: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines. Decision based on Non-MTUS Citation Official Disability Guidelines (ODG).

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Benzodiazepines Page(s): 24.

Decision rationale: MTUS states that benzodiazepine (ie Xanax) is "Not recommended for long-term use because long-term efficacy is unproven and there is a risk of dependence. Most guidelines limit use to 4 weeks. Their range of action includes sedative/hypnotic, anxiolytic, anticonvulsant, and muscle relaxant. Chronic benzodiazepines are the treatment of choice in very few conditions. Tolerance to hypnotic effects develops rapidly. Tolerance to anxiolytic effects occurs within months and long-term use may actually increase anxiety. A more appropriate treatment for anxiety disorder is an antidepressant. Tolerance to anticonvulsant and muscle relaxant effects occurs within weeks." Medical records indicate that the patient has been on Xanax in the past far exceeding MTUS recommendations. The medical record does not provide any extenuating circumstances to recommend exceeding the guideline recommendations. Additionally, no documentation as to if a trial of antidepressants was initiated and the outcome of this trial. As such, the request is not medically necessary.

Norco 10/325 mg #90: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Opioids.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Opioids Page(s): 74-96. Decision based on Non-MTUS Citation Official Disability Guidelines (ODG) Neck and Upper Back (Acute and Chronic), Low Back - Lumbar & Thoracic (Acute & Chronic), Opioids, Pain.

Decision rationale: ODG does not recommend the use of opioids for neck, low back, and shoulder pain "except for short use for severe cases, not to exceed 2 weeks." The patient has exceeded the 2 week recommended treatment length for opioid usage. MTUS does not discourage use of opioids past 2 weeks, but does state that "ongoing review and documentation of pain relief, functional status, appropriate medication use, and side effects. Pain assessment should include current pain; the least reported pain over the period since last assessment; average pain; intensity of pain after taking the opioid; how long it takes for pain relief; and how long pain relief lasts. Satisfactory response to treatment may be indicated by the patient's decreased pain, increased level of function, or improved quality of life." The treating physician does not fully document the least reported pain over the period since last assessment, intensity of pain after taking opioid, pain relief, increased level of function, or improved quality of life. Additionally, medical documents indicate that the patient has been on Norco in excess of the recommended 2-week limit. As such, the request for Norco is not medically necessary.

Oxycodone IR 10mg #40: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Opioids.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Opioids Page(s): 74-96. Decision based on Non-MTUS Citation Official Disability Guidelines (ODG) Neck and Upper Back (Acute and Chronic), Low Back - Lumbar & Thoracic (Acute & Chronic), Opioids, Pain.

Decision rationale: ODG does not recommend the use of opioids for neck, low back, and shoulder pain "except for short use for severe cases, not to exceed 2 weeks." The patient has exceeded the 2 week recommended treatment length for opioid usage. MTUS does not discourage use of opioids past 2 weeks, but does state that "ongoing review and documentation of pain relief, functional status, appropriate medication use, and side effects. Pain assessment should include current pain; the least reported pain over the period since last assessment; average pain; intensity of pain after taking the opioid; how long it takes for pain relief; and how long pain relief lasts. Satisfactory response to treatment may be indicated by the patient's decreased pain, increased level of function, or improved quality of life." The treating physician does not fully document the least reported pain over the period since last assessment, intensity of pain after taking opioid, pain relief, increased level of function, or improved quality of life. Additionally, medical documents indicate that the patient has been on an opioid in excess of the recommended 2-week limit. As such, the request for Oxycodone is not medically necessary.

TPI's for (L) lumbar deep fascia: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Trigger Point Injections.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Trigger point injections Page(s): 122.

Decision rationale: MTUS states that Trigger Point Injections are "Recommended only for myofascial pain syndrome as indicated below, with limited lasting value. Not recommended for radicular pain." Further states that "trigger point is a discrete focal tenderness located in a palpable taut band of skeletal muscle, which produces a local twitch in response to stimulus to the band. For fibromyalgia syndrome, trigger points injections have not been proven effective." MTUS lists the criteria for Trigger Points: (1) Documentation of circumscribed trigger points with evidence upon palpation of a twitch response as well as referred pain; (2) Symptoms have persisted for more than three months; (3) Medical management therapies such as ongoing stretching exercises, physical therapy, NSAIDs and muscle relaxants have failed to control pain; (4) Radiculopathy is not present (by exam, imaging, or neuro-testing); (5) Not more than 3-4 injections per session; (6) No repeat injections unless a greater than 50% pain relief is obtained for six weeks after an injection and there is documented evidence of functional improvement; (7) Frequency should not be at an interval less than two months; (8) Trigger point injections with any substance (e.g., saline or glucose) other than local anesthetic with or without steroid are not recommended. The medical documents do meet some criteria for trigger point injections per MTUS. MTUS specifically states that radiculopathy should not be present by exam, imaging, or neuro-testing. However, subjective complaints of radiculopathy are present on numerous treatment notes. As such, the request for trigger point injections is not medically necessary.