

Case Number:	CM15-0042926		
Date Assigned:	03/13/2015	Date of Injury:	11/08/2000
Decision Date:	04/17/2015	UR Denial Date:	02/14/2015
Priority:	Standard	Application Received:	03/06/2015

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:

State(s) of Licensure: Indiana

Certification(s)/Specialty: Preventive Medicine, Occupational Medicine

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The 39-year-old male injured worker suffered an industrial injury on 11/8/2000. The diagnoses were intractable lumbar pain, lumbar radiculopathy, and chronic cervical pain with radiculitis, bilateral knee tendinosis, depression, and anxiety. The treatments were medications. The treating provider reported significant back pain with neck pain and knee pain. On exam there was impaired gait using a walker with significant guarding and tenderness over the cervical and lumbar spine. The requested treatments were: 1. Oxycodone 10mg #60. 2. Fentanyl Patch #10. 3. Zanaflex 4mg #60.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

Oxycodone 10mg #60: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Opioids Page(s): 74-97.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Opioids Page(s): 74-96. Decision based on Non-MTUS Citation Official Disability Guidelines (ODG) Low Back - Lumbar & Thoracic (Acute & Chronic) and Pain, Opioids.

Decision rationale: Oxycodone is the generic version of Oxycotin, which is a pure opioid agonist. ODG does not recommend the use of opioids for low back pain "except for short use for severe cases, not to exceed 2 weeks." The patient has exceeded the 2 week recommended treatment length for opioid usage. MTUS does not discourage use of opioids past 2 weeks, but does state that "ongoing review and documentation of pain relief, functional status, appropriate medication use, and side effects. Pain assessment should include: current pain; the least reported pain over the period since last assessment; average pain; intensity of pain after taking the opioid; how long it takes for pain relief; and how long pain relief lasts. Satisfactory response to treatment may be indicated by the patient's decreased pain, increased level of function, or improved quality of life." The treating physician does not fully document the least reported pain over the period since last assessment, intensity of pain after taking opioid, pain relief, increased level of function, or improved quality of life. As such, the question for Oxycodone is not medically necessary.

Fentanyl Patch #10: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Opioids Page(s): 74-97.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Fentanyl Page(s): 44. Decision based on Non-MTUS Citation Official Disability Guidelines (ODG) Pain, Opioids, Specific drug list.

Decision rationale: CA MTUS states and ODG agrees: "Not recommended as a first-line therapy. Duragesic is the trade name of a fentanyl transdermal therapeutic system, which releases fentanyl, a potent opioid, slowly through the skin. The FDA-approved product labeling states that Duragesic is indicated in the management of chronic pain in patients who require continuous opioid analgesia for pain that cannot be managed by other means." ODG does not recommend the use of opioids "except for short use for severe cases, not to exceed 2 weeks." The patient has exceeded the 2 week recommended treatment length for opioid usage. MTUS does not discourage use of opioids past 2 weeks, but does state that "ongoing review and documentation of pain relief, functional status, appropriate medication use, and side effects. Pain assessment should include current pain; the least reported pain over the period since last assessment; average pain; intensity of pain after taking the opioid; how long it takes for pain relief; and how long pain relief lasts. Satisfactory response to treatment may be indicated by the patient's decreased pain, increased level of function, or improved quality of life." The treating physician does not include pain assessments such as current, least, and average. With the multiple pain medications, there is serious risk of opioid dependence. Therefore, the request for Fentanyl Patch #10 is not medically necessary.

Zanaflex 4mg #60: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Muscle relaxants (for pain) Page(s): 63-66.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Muscle relaxants Page(s): 63-67.

Decision rationale: Zanaflex is the brand name version of tizanidine, which is a muscle relaxant. MTUS states concerning muscle relaxants "Recommend non-sedating muscle relaxants with caution as a second-line option for short-term treatment of acute exacerbations in patients with chronic LBP. (Chou, 2007) (Mens, 2005) (VanTulder, 1998) (van Tulder, 2003) (van Tulder, 2006) (Schnitzer, 2004) (See, 2008) Muscle relaxants may be effective in reducing pain and muscle tension, and increasing mobility. However, in most LBP cases, they show no benefit beyond NSAIDs in pain and overall improvement. In addition, there is no additional benefit shown in combination with NSAIDs. Efficacy appears to diminish over time, and prolonged use of some medications in this class may lead to dependence. (Homik, 2004) Sedation is the most commonly reported adverse effect of muscle relaxant medications. These drugs should be used with caution in patients driving motor vehicles or operating heavy machinery. Drugs with the most limited published evidence in terms of clinical effectiveness include chlorzoxazone, methocarbamol, dantrolene and baclofen. (Chou, 2004) According to a recent review in American Family Physician, skeletal muscle relaxants are the most widely prescribed drug class for musculoskeletal conditions (18.5% of prescriptions), and the most commonly prescribed antispasmodic agents are carisoprodol, cyclobenzaprine, metaxalone, and methocarbamol, but despite their popularity, skeletal muscle relaxants should not be the primary drug class of choice for musculoskeletal conditions. (See2, 2008)" MTUS further states, "Tizanidine (Zanaflex, generic available) is a centrally acting alpha2-adrenergic agonist that is FDA approved for management of spasticity; unlabeled use for low back pain. (Malanga, 2008) Eight studies have demonstrated efficacy for low back pain. (Chou, 2007) One study (conducted only in females) demonstrated a significant decrease in pain associated with chronic myofascial pain syndrome and the authors recommended its use as a first line option to treat myofascial pain. (Malanga, 2002) May also provide benefit as an adjunct treatment for fibromyalgia. (ICSI, 2007)" It is unclear how long the employee has been on this medication and what relief he is getting, and how this medication fits into his overall pain control plan. Therefore, the request for Zanaflex is not medically necessary.