

<b>Case Number:</b>	CM15-0043671		
<b>Date Assigned:</b>	03/13/2015	<b>Date of Injury:</b>	01/14/2014
<b>Decision Date:</b>	04/24/2015	<b>UR Denial Date:</b>	02/06/2015
<b>Priority:</b>	Standard	<b>Application Received:</b>	03/09/2015

### HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:  
State(s) of Licensure: North Carolina  
Certification(s)/Specialty: Family Practice

### CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The injured worker is a 45 year old male, who sustained an industrial injury on January 14, 2014. He has reported left shoulder pain and right thumb and hand pain. Diagnoses have included left rotator cuff tear, and crushing injury of the hand and fingers. Treatment to date has included medications, physical therapy, acupuncture, left shoulder surgery, and imaging studies. A progress note dated December 15, 2014 indicates a chief complaint of left shoulder pain. The treating physician documented a plan of care that included medications and a sleep study referral.

### IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

**Compound: Panthenol, Bupivacaine, Gabapentin, Amitriptyline, 2140gm x30 days (12/23/2014): Upheld**

**Claims Administrator guideline:** Decision based on MTUS Chronic Pain Treatment Guidelines Topical Analgesics Page(s): 111-113.

**MAXIMUS guideline:** Decision based on MTUS Chronic Pain Treatment Guidelines topical analgesics Page(s): 111-113.

**Decision rationale:** The California chronic pain medical treatment guidelines section on topical analgesics states: Recommended as an option as indicated below. Largely experimental in use with few randomized controlled trials to determine efficacy or safety. Primarily recommended for neuropathic pain when trials of antidepressants and anticonvulsants have failed. (Namaka, 2004) These agents are applied locally to painful areas with advantages that include lack of systemic side effects, absence of drug interactions, and no need to titrate. (Colombo, 2006) Many agents are compounded as monotherapy or in combination for pain control (including NSAIDs, opioids, capsaicin, local anesthetics, antidepressants, glutamate receptor antagonists, adrenergic receptor agonist, adenosine, cannabinoids, cholinergic receptor agonists, agonists, prostanoids, bradykinin, adenosine triphosphate, biogenic amines, and nerve growth factor). (Argoff, 2006) There is little to no research to support the use of many of these agents. Any compounded product that contains at least one drug (or drug class) that is not recommended is not recommended. The requested medication contains multiple ingredients, which are not indicated per the California MTUS for topical analgesic use. Therefore, the request is not certified and is not medically necessary.

**Compound: Flurbiprofen, Beclofen, Dexamethasone, Menthol, Camphor, capsaicin, 310gm, x 30 days (12/23/2014): Upheld**

**Claims Administrator guideline:** Decision based on MTUS Chronic Pain Treatment Guidelines Topical Analgesics Page(s): 111-113.

**MAXIMUS guideline:** Decision based on MTUS Chronic Pain Treatment Guidelines topical analgesics Page(s): 111-113.

**Decision rationale:** The California chronic pain medical treatment guidelines section on topical analgesics states: Recommended as an option as indicated below. Largely experimental in use with few randomized controlled trials to determine efficacy or safety. Primarily recommended for neuropathic pain when trials of antidepressants and anticonvulsants have failed. (Namaka, 2004) These agents are applied locally to painful areas with advantages that include lack of systemic side effects, absence of drug interactions, and no need to titrate. (Colombo, 2006) Many agents are compounded as monotherapy or in combination for pain control (including NSAIDs, opioids, capsaicin, local anesthetics, antidepressants, glutamate receptor antagonists, adrenergic receptor agonist, adenosine, cannabinoids, cholinergic receptor agonists, agonists, prostanoids, bradykinin, adenosine triphosphate, biogenic amines, and nerve growth factor). (Argoff, 2006) There is little to no research to support the use of many of these agents. Any compounded product that contains at least one drug (or drug class) that is not recommended is not recommended. The requested medication contains multiple ingredients, which are not indicated per the California MTUS for topical analgesic use. Therefore, the request is not certified and is not medically necessary.