

Case Number:	CM15-0043969		
Date Assigned:	03/13/2015	Date of Injury:	10/19/2002
Decision Date:	04/22/2015	UR Denial Date:	03/03/2015
Priority:	Standard	Application Received:	03/09/2015

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:
 State(s) of Licensure: New Jersey, Michigan, California
 Certification(s)/Specialty: Neurology, Neuromuscular Medicine

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The injured worker is a 43 year old female, who sustained an industrial injury on October 19, 2002. The diagnoses have included subacromial bursitis, cervical sprain and rule out neuropathy. Treatment to date was not included in the physician's progress report. Currently, the injured worker complains of neck, and right shoulder pain. In a progress note dated February 13, 2015, the treating provider reports examination of right shoulder revealed decreased range of motion and tenderness to palpation.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

MRI Scan of the right shoulder: Upheld

Claims Administrator guideline: Decision based on MTUS ACOEM Chapter 9 Shoulder Complaints.

MAXIMUS guideline: Decision based on MTUS ACOEM Chapter 9 Shoulder Complaints
 Page(s): 209.

Decision rationale: According to MTUS guidelines, MRI of the shoulder is indicated in case of tumor, infection, ligament instability and rotator cuff injury. There is no clinical evidence or documentation of one of the above diagnosis. Therefore MRI of the right shoulder is not medically necessary.

EMG of the bilateral upper extremities: Upheld

Claims Administrator guideline: Decision based on MTUS ACOEM Chapter 9 Shoulder Complaints.

MAXIMUS guideline: Decision based on MTUS ACOEM Chapter 11 Forearm, Wrist, and Hand Complaints Page(s): 178.

Decision rationale: According to MTUS guidelines (MTUS page 303 from ACOEM guidelines), “Electromyography (EMG), including H-reflex tests, may be useful to identify subtle, focal neurologic dysfunction in patients with low back symptoms lasting more than three or four weeks.” EMG has excellent ability to identify abnormalities related to disc protrusion (MTUS page 304 from ACOEM guidelines). According to MTUS guidelines, needle EMG study helps identify subtle neurological focal dysfunction in patients with neck and arm symptoms. “When the neurologic examination is less clear, however, further physiologic evidence of nerve dysfunction can be obtained before ordering an imaging study
Electromyography (EMG), and nerve conduction velocities (NCV), including H-reflex tests, may help identify subtle focal neurologic dysfunction in patients with neck or arm symptoms, or both, lasting more than three or four weeks” (page 178). EMG is indicated to clarify nerve dysfunction in case of suspected disc herniation (page 182). EMG is useful to identify physiological insult and anatomical defect in case of neck pain (page 179). The patient developed chronic neck pain without recent evidence of radicular pain and no recent clear justification for the need of an EMG. Therefore, the request for EMG of bilateral upper extremities is not medically necessary.

Tramadol 50mg #120: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Opioids.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Tramadol Page(s): 113.

Decision rationale: According to MTUS guidelines, Ultram (Tramadol) is a synthetic opioid indicated for the pain management but not recommended as a first line oral analgesic. In addition and according to MTUS guidelines, ongoing use of opioids should follow specific rules: (a) Prescriptions from a single practitioner taken as directed, and all prescriptions from a single pharmacy. (b) The lowest possible dose should be prescribed to improve pain and function. (c) Office: Ongoing review and documentation of pain relief, functional status, appropriate medication use, and side effects. Pain assessment should include: current pain; the least reported pain over the period since last assessment; average pain; intensity of pain after taking the opioid;

how long it takes for pain relief; and how long pain relief lasts. Satisfactory response to treatment may be indicated by the patient's decreased pain, increased level of function, or improved quality of life. Information from family members or other caregivers should be considered in determining the patient's response to treatment. The 4 A's for Ongoing Monitoring: Four domains have been proposed as most relevant for ongoing monitoring of chronic pain patients on opioids: pain relief, side effects, physical and psychosocial functioning, and the occurrence of any potentially aberrant (or non adherent) drug-related behaviors. These domains have been summarized as the "4 A's" (analgesia, activities of daily living, adverse side effects, and aberrant drug taking behaviors). The monitoring of these outcomes over time should affect therapeutic decisions and provide a framework. There is no clear documentation of pain and functional improvement with previous use of the Tramadol. There is no clear documentation of continuous documentation of patient compliance with her medications. There is no documentation of the medical necessity of Tramadol over NSAID. Therefore, the prescription of Tramadol 50 mg #120 is not medically necessary.